



PHOENIX

ENVIRONMENTAL SCIENCES

Level 1 subterranean fauna assessment for the Beyondie Sulphate of Potash Project

Prepared for Kalium Lakes Potash Pty Ltd

March 2018

Final Report



Level 1 subterranean fauna assessment for the Beyondie Sulphate of Potash Project

Prepared for Kalium Lakes Potash Pty Ltd

Final Report

Author/s: Volker Framenau, Jarrad Clark

Reviewer: Karen Crews

Date: 14 March 2018

Submitted to: Gavin Edwards, Phil Scott, Brett Hazelden

Chain of authorship and review			
Name	Task	Version	Date
V.W. Framenau	Draft for client comments	1.0	6 June 2017
J. Clark	Final submitted to client	2.0	14 March 2018
	Choose an item.		

NOTE ON REPORT CURRENCY

This report was prepared prior to the completion of a detailed H3 hydrological assessment for the Beyondie Sulphate of Potash Project (Advisian 2017a) and therefore some hydrological information in this report may not be current. Updated hydrological information and interpretation of its relevance for subterranean fauna will be presented in the subsequent level 2 subterranean fauna survey report for the Project.

©Phoenix Environmental Sciences Pty Ltd 2018

The use of this report is solely for the Client for the purpose in which it was prepared. Phoenix Environmental Sciences accepts no responsibility for use beyond this purpose.

All rights are reserved and no part of this report may be reproduced or copied in any form without the written permission of Phoenix Environmental Sciences or the Client.

[Phoenix Environmental Sciences Pty Ltd](#)

1/511 Wanneroo Rd BALCATTA WA 6021

P: 08 96323 5410

E: admin@phoenixenv.com.au

Project code: 1148-TM-KAL-ECO

Contents

CONTENTS.....	I
LIST OF FIGURES.....	II
LIST OF TABLES.....	II
LIST OF APPENDICES	II
EXECUTIVE SUMMARY	III
1 INTRODUCTION.....	5
1.1 Background	5
1.2 Survey objectives and scope of works	7
1.3 Study area	7
2 LEGISLATIVE CONTEXT	7
2.1 Commonwealth.....	7
2.2 State	8
3 EXISTING ENVIRONMENT.....	10
3.1 Interim Biogeographic Regionalisation of Australia	10
3.2 Climate and weather.....	12
3.3 Geology	13
3.3.1 Brine extraction area.....	13
3.3.2 Freshwater development envelope.....	13
3.4 Hydrology.....	15
3.4.1 Brine extraction area.....	15
3.4.2 Freshwater development envelope.....	15
3.5 Biological context.....	16
3.5.1 Stygofauna	16
3.5.2 Troglofauna	17
3.5.3 Threatening processes	17
3.5.4 Calcretes as subterranean fauna habitat	18
3.5.5 Salinity.....	19
4 METHODS.....	20
4.1 Desktop review	20
4.1.1 Database searches	20
4.1.2 Literature review.....	20
4.2 Field survey	21
4.2.1 Survey.....	21
4.2.2 Fauna sampling	23
4.2.3 Bore and water quality data	23
4.2.4 Morphological species identifications	23
4.2.5 Molecular identifications	23
4.3 Project personnel.....	24
5 RESULTS	25

5.1	Desktop review	25
5.1.1	Database searches	25
5.1.2	Threatened and Priority species or ecological communities	25
5.1.3	Literature review	25
5.2	Field survey	28
5.2.1	Water quality	28
5.2.2	Subterranean fauna	29
5.3	Survey limitations	34
6	DISCUSSION.....	35
6.1	Recommendations	36
7	REFERENCES.....	37

List of Figures

Figure 1-1	Location of the Beyondie Potash Project and study area for the Level 1 subterranean fauna assessment.....	6
Figure 3-1	Location of the Beyondie Sulphate of Potash Project in relation to IBRA regions and subregions.....	11
Figure 3-2	Average monthly temperatures and rainfall for Newman (BoM 2017)	12
Figure 3-3	Surface geology of the Beyondie Potash Project	14
Figure 4-1	Location of bores sampled during field survey	22
Figure 5-1	Subterranean fauna records from the desktop review	27
Figure 5-2	Subterranean fauna collected during field survey.....	31

List of Tables

Table 4-1	Selected aquatic invertebrate surveys of salt lakes in Western Australia.....	21
Table 4-2	Location and details of bores sampled during field survey	21
Table 4-3	Project team.....	24
Table 5-1	Subterranean fauna recorded in the desktop review.....	26
Table 5-2	Water quality data for sampled bores.....	28
Table 5-3	Potential stygofauna fauna collected during field survey	29
Table 5-4	Potential troglafauna fauna collected during field survey	30
Table 5-5	Survey limitations based on EPA Technical Guide: Terrestrial fauna surveys (EPA 2016e)	34

List of Appendices

Appendix 1	Conservation significant subterranean fauna in Western Australia (DPaW 2017)
Appendix 2	Subterranean Threatened Ecological Communities in Western Australia (DPaW 2016a)
Appendix 3	Priority Ecological Communities (DPaW 2016b)

EXECUTIVE SUMMARY

Kalium Lakes Potash Pty Ltd (Kalium) proposes to develop the Beyondie Sulphate of Potash Project (the Project). The Project is located approximately 150 km south-southeast of Newman with access to the Great Northern Highway at Kumarina approximately 65 km to the east. The Project spans the border between the Little Sandy Desert and Gascoyne bioregions. Kalium proposes to produce Sulphate of Potash (SOP) by abstraction of hypersaline groundwater brine containing the dissolved product, which is extracted by water evaporation.

Phoenix Environmental Sciences Pty Ltd (Phoenix) was commissioned to conduct a level 1 subterranean fauna assessment for the Project to support environmental approvals. The study area for this assessment is loosely defined as 1) the exploration leases E69/3309, E69/3346, E69/3347, E69/3351 and E69/3352; 2) a development envelope for the provision of fresh water ('Freshwater Development Envelope') and regional sampling bores sampled outside these areas.

The objective of this level 1 subterranean fauna survey was to assess the likelihood of subterranean fauna to occur in the study area by reviewing relevant databases, available technical reports, literature and spatial data, to determine whether a pilot subterranean fauna survey was required. The pilot survey was conducted from 29–31 March 2017 targeting a total of 15 bores and wells, seven within the study area and eight considered regional reference sites.

The methodology of this assessment complied with the Environmental Protection Authority's (EPA's) requirements for an environmental impact assessment (EIA) of subterranean invertebrates as outlined in the *Environmental Factor Guideline: Subterranean fauna* and *Technical Guidance: Sampling methods for subterranean fauna*.

Subterranean fauna are animals, predominantly invertebrates, which have evolved to live underground to escape harsh environmental conditions such as extreme heat and dryness of exposed environments. They are classified into two types:

- troglofauna – animals that live in air-filled subterranean networks
- stygofauna – animals that live in water-filled subterranean networks.

Habitats likely to support troglofauna are karstic limestone, channel iron deposits (CID; in particular pisolite in inverted landscape geomorphology), groundwater calcretes above the water table, alluvium/colluvium in valley-fill settings, banded iron formations (BIF) and weathered and fractured sandstone. Stygofauna are likely where there are groundwater voids present, for example in karst limestone, calcretes, alluvial formations and fractured rock.

The Project is located just off the north-western edge of the Yilgarn Craton and just off the south-western edge of the North-west Officer Basin. Beyondie Lakes, Ten Mile Lake and Lake Sunshine are developed within a palaeochannel system within the Little Sandy Desert north of the Yilgarn Craton, underlain by bedrock of the Sunbeam Group, consisting mostly of sedimentary sandstones, siltstones, conglomerates and shales. Within the lakes, Quaternary low porosity lacustrine deposits are mainly clay, mud and silt which are hypersaline and commonly gypsiferous. Surrounding the lakes are expanses of Quaternary Aeolian sand and sand sheets. Longitudinal (seif), chain and net dunes are abundant. Two large calcrete deposits present and are reflected in two surface drainage lines, i.e. Nanyeriny Creek flowing into the Beyondie Lakes from the west and 477 Creek flowing into Ten Mile Lake from the south.

Four main aquifers exist in the study area, including upper unconsolidated alluvial and colluvial sediments (<15 m deep), areas of saturated calcrete, basal palaeochannel sands and areas of weathered or fractured bedrock.

Database searches identified seven stygofauna species within approximately 100 km of the study area. Three of these are only known from single localities: *Billibathynella ilgarariensis* and *Brevisomabathynella magna* (both Parabathynellidae) and *Pilbaracyclops frustratio* (Cyclopidae). All others are widespread species. No troglifauna species were identified in the desktop review.

The desktop study therefore concluded that within the lake containing palaeochannels, low porosity lacustrine deposits (of clay, mud and silt), which are hypersaline (70,000 to over 300,000 mg/L TDS) have a low prospectivity for subterranean fauna. And conversely, that the alluvial freshwater aquifers targeted by the Freshwater Development Envelope had high potential for stygofauna generally (and troglifauna above the water table, which will not be impacted), and especially in calcrete formations.

The pilot survey recovered two taxa of stygofauna, syncarid crustaceans (order Bathynellacea) and amphipods (family Paramelitidae) from Garden Well and Tupee Well. Molecular analysis of the Garden Well specimens identified two species, an amphipod in the family Paramelitidae (AMP036) and a syncarid in the family Parabathynellidae (BAP028) that have not previously been recorded elsewhere. Amphipod specimens from Tupee Well were not able to be sequenced to test for conspecificity with the Garden Well species. Two species of troglifauna were also recovered; an unidentified spider (Garden Well) and a juvenile meenoplid bug (Hemiptera) (David's Well). All remaining species were found to be aquatic surface taxa, stygophiles or edaphic (soil) fauna.

As expected the subterranean fauna records were from bores in alluvial deposits at considerable distance west of the Beyondie Lakes and Ten Mile Lake, no sampling was undertaken at Lake Sunshine. Garden Well is within the Freshwater Development Envelope, Tupee Well is not.

At this stage the Nanyerinni Creek calcrete formation has not been sampled but is not expected to be impacted by either brine extraction or freshwater abstraction for the Project.

The extremely large calcrete formation (15 km x 7.5 km) associated with Ten Mile Lake has only been sampled with a single bore. The lacustrine, hypersaline aquifers associated with Ten Mile Lake and Lake Sunshine are considered unlikely to support stygofauna and troglifauna.

The pilot study has resulted in records of both troglifauna and stygofauna in the alluvial aquifers of the Freshwater Development Envelope, west of the Beyondie and Ten Mile Lake systems. This result therefore triggers the need for a level 2 subterranean survey for the Project consistent with EPA guidelines. Limited groundwater drawdown is proposed for the aquifers and consequently, it is unlikely that groundwater abstraction within the Freshwater Development Envelope will have significant impacts on troglifauna. It is therefore recommended that future surveys focus on stygofauna sampling in the aquifers of the Freshwater Development Envelope.

1 INTRODUCTION

Phoenix Environmental Sciences Pty Ltd (Phoenix) was commissioned by Preston Consulting Pty Ltd (Preston) on behalf of Kalium Lakes Potash Pty Ltd (Kalium) to conduct a level 1 subterranean fauna assessment for the Beyondie Sulphate of Potash Project (the Project). The Project is located approximately 150 km south-southeast of Newman, with access to the Great Northern Highway at Kumarina approximately 65 km to the east (Figure 1-1). The project area spans the border between the Great Sandy Desert and Gascoyne bioregions.

1.1 BACKGROUND

The Beyondie Lakes consist of a western freshwater marsh (outside the study area) connected to two circular salt playas in line in the east. Ten Mile Lake is a large salt playa located about 6 km to the south. Several claypans are located around the lakes. The Beyondie Lakes salt playas connect with Ten Mile Lake during extreme inundation events.

Kalium is seeking to develop a sub-surface brine deposit to produce 150 ktpa Sulphate of Potash (SOP) product via an evaporation and processing operation. The Project will entail the extraction of hypersaline groundwater that will be pumped to ponds where the salts will be concentrated, and different salts extracted and purified sequentially.

The Project is based on the extraction of hypersaline groundwater from beneath the salt lake systems. It particularly targets the basal palaeochannel aquifers that are likely to have a significantly higher hydraulic conductivity and specific yield than the other layers in the profile. However, brine abstraction by superficial channels on the salt lake also forms part of the Project.


A number of piezometers will be established around the test bores selected for extended pump testing. These will serve to monitor the propagation of drawdown within the aquifer. It is anticipated that the target aquifer will behave as a confined or semi-confined aquifer. A trial project will be undertaken that involves the pumping of 4–6 test bores to supply trial ponds (Advisian 2017a).

Considering that the basal aquifer has a 30–40 m confining head, the drop-in water levels is expected to leave the aquifer system in a confined state after a year of pumping. Downward leakage from the clays and the silts/sands above the main aquifer would likely occur, but with a drop-in water levels far less than in the basal sand aquifer due to low porosity and high clay content and therefore, the impact on the aquifers above the clays would be expected to be very limited.


Calcrete aquifers to the south and west of Ten Mile Lake are thought contain fresh water, suitable for processing requirements, at this stage only the calcrete associated with Ten Mile Lake is being targeted (see Freshwater Development Envelope in Figure 1-1). Proposed abstraction here is expected to be no more than 1.5 GL per annum (Preston Consulting 2017).

Figure 1–1
Location of the Beyondie Potash Project and study area for the Level 1 subterranean fauna assessment

-  Freshwater Development Envelope
-  Tenement E69/3309
-  Tenement E69/3346
-  Tenement E69/3347
-  Tenement E69/3351
-  Tenement E69/3352
-  Road
-  Minor road
-  Major creeks and rivers
-  Lake



Client: Kalium Lakes Ltd
 Project: Beyondie Sulphate of Potash Project
 Author: AL
 Date: 12-Jan-18
 Coordinate System: GDA 1994 MGA Zone 51
 Projection: Transverse Mercator
 Datum: GDA 1994



Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community

1.2 SURVEY OBJECTIVES AND SCOPE OF WORKS

The objective of the level 1 subterranean fauna assessment was to identify the likelihood that subterranean fauna are present in the study area, with particular reference to the abstraction of the potassium hosting brine (hypersaline water) from the underlying lake aquifers and sourcing of fresh process water from regional alluvial deposits.

The scope of work was as follows:

- conduct a desktop review of available technical reports, relevant databases and spatial data to assess the potential for presence of subterranean fauna in the study area
- conduct a pilot survey for subterranean fauna within prospective geologies within the study area
- prepare a technical report detailing the desktop and field results that considers separately the communities and species associated with the process water from the alluvial aquifers and those associated with Ten Mile Lake and Lake Sunshine, including recommendations on additional sampling, if required.

The assessment methodology complied with the Environmental Protection Authority's (EPA's) requirements for an environmental impact assessment (EIA) of subterranean invertebrates as outlined in Environmental Factor Guideline: Subterranean fauna (EPA 2016a) and Technical Guidance: Sampling methods for subterranean fauna (EPA 2016c).

1.3 STUDY AREA

The study area for the level 1 subterranean fauna assessment was loosely defined as 1) the exploration leases E69/3309, E69/3346, E69/3347, E69/3351 and E69/3352, and 2) the Freshwater Development Envelope (Preston Consulting 2017) (Figure 1-1), but regional surveys in particular along the Nanyeriny Creek system in the west of the study area complemented the assessment.

2 LEGISLATIVE CONTEXT

The protection of fauna in Western Australia (WA) is principally governed by three Acts:

- Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act)
- *Wildlife Conservation Act 1950* (WC Act)
- *Environmental Protection Act 1986* (EP Act).

The WA *Biodiversity Conservation Act 2016* (BC Act) will eventually replace the WC Act; however, the provisions in the BC Act pertaining to the listing of flora and fauna cannot be brought into effect until the necessary Biodiversity Conservation Regulations have been made.

2.1 COMMONWEALTH

Under the EPBC Act, actions that have, or are likely to have, a significant impact on a matter of national environmental significance (NES), require approval from the Australian Government Minister for the Environment.

The EPBC Act provides for the listing of Threatened fauna, flora and ecological communities as matters of NES. Conservation categories applicable to Threatened fauna, flora and ecological communities under the EPBC Act are:

- Extinct (EX)¹ – there is no reasonable doubt that the last individual has died
- Extinct in the Wild (EW) – taxa known to survive only in captivity
- Critically Endangered (CR) – taxa facing an extremely high risk of extinction in the wild in the immediate future
- Endangered (EN) – taxa facing a very high risk of extinction in the wild in the near future
- Vulnerable (VU) – taxa facing a high risk of extinction in the wild in the medium-term
- Conservation Dependent¹ – taxa whose survival depends upon ongoing conservation measures; without these measures, a conservation dependent taxon would be classified as Vulnerable or more severely threatened.

2.2 STATE

In WA, the WC Act provides for the listing of native fauna (Threatened Fauna) species which are under identifiable threat of extinction². Threatened Fauna are assigned to one of seven categories under the WC Act (Western Australian Government 2015):

- Schedule 1 – fauna that is rare or is likely to become extinct as critically endangered (CR) fauna
- Schedule 2 – fauna that is rare or is likely to become extinct as endangered (EN) fauna
- Schedule 3 – fauna that is rare or is likely to become extinct as vulnerable (VU) fauna
- Schedule 4 – fauna presumed to be extinct (EX)
- Schedule 5 – Migratory birds protected under an international agreement
- Schedule 6 – fauna that is of special conservation need as conservation dependent fauna
- Schedule 7 – other specially protected (SP) fauna.

Assessments for listing of fauna are based on the International Union for Conservation of Nature (IUCN) threat categories.

The Department of Biodiversity, Conservation and Attractions (DBCA) administers the WC Act and also maintains a non-statutory list of Priority fauna species. Priority species are still considered to be of conservation significance – that is they may be rare or threatened – but cannot be considered for listing under the WC Act until there is adequate understanding of their threat levels. Species on the Priority fauna lists are assigned to one of five Priority (P) categories, P1 (highest) – P5 (lowest), based on level of knowledge/concern.

¹ Species listed as Extinct and Conservation Dependent are not matters of NES and therefore do not trigger the EPBC Act.

² This function of the WC Act will be replaced by the BC Act when the relevant BC Act regulations come into effect.

A total of 23 troglofauna and 20 stygofauna species are currently listed as either Threatened or Priority in WA, with the majority from the Pilbara and Carnarvon Interim Biogeographic Regionalisation of Australia (IBRA) regions (Appendix 1).

The Minister for Environment may also list ecological communities which are at risk of becoming destroyed as 'Threatened'³. DBCA maintains a list of ministerially-endorsed Threatened Ecological Communities (TECs) (DPaW 2016a) as well as a non-statutory list of Priority Ecological Communities (PECs) (DPaW 2016b) which are also assigned to one of five categories.

Nine subterranean TECs (Appendix 2) and 86 subterranean PECs (Appendix 3) are listed from WA. Of the 86 PECs, 77 are stygofauna communities in the groundwater of calcretes of the Yilgarn Craton in the Midwest and northern Goldfields regions (i.e. Cooper *et al.* 2008; Guzik *et al.* 2008; Humphreys *et al.* 2009).

³ The BC Act will allow for the listing of TECs when the relevant BC Act regulations come into effect.

3 EXISTING ENVIRONMENT

3.1 INTERIM BIOGEOGRAPHIC REGIONALISATION OF AUSTRALIA

The IBRA defines 'bioregions' as large land areas characterised by broad, landscape-scale natural features and environmental processes that influence the functions of entire ecosystems (Department of the Environment 2014b; Thackway & Cresswell 1995). They categorise the large-scale geophysical patterns that occur across the Australian continent that are linked to fauna and flora assemblages and processes at the ecosystem scale (Thackway & Cresswell 1995).

Western Australia contains 26 IBRA bioregions and 53 subregions. The study area is situated on the border to two bioregions; the Gascoyne bioregion and Little Sandy Desert bioregion (Figure 3-1); At the junction of the Augustus subregion (GAS3) of the Gascoyne bioregion and Trainor subregion (LSD2) of the Little Sandy Desert bioregion (Figure 3-1).

The Trainor subregion (LSD2) contains the target potassium hosting aquifers and is characterised by (Cowan & Kendrick 2001):

- red centre desert on Neoproterozoic sedimentary basement (Officer Basin)
- red Quaternary dune fields with abrupt Proterozoic sandstone ranges of Bangemall Basin
- shrub-steppe of acacias, *Aluta maisonneuvei* and grevilleas over *Triodia schinzii* on sandy surfaces
- sparse shrub-steppe over *Triodia basedowii* on stony hills
- eucalypt and coolabah communities and bunch grasses on alluvial deposits and drainage lines associated with ranges
- arid climate with episodic summer rainfall.

The Augustus subregion (GAS3) contains the target freshwater processing aquifers and is characterised by (Desmond *et al.* 2001):


- low Proterozoic sedimentary and granite ranges divided by flat broad valleys
- mulga woodland with *Triodia* on shallow stony loams on rises with mulga parkland on shallow earthy loams over hardpan on the plains
- extensive areas of alluvial deposits
- calcrete aquifers of the Carnegie drainage system
- desert climate with bimodal rainfall.

With respect to subterranean fauna, the alluvial deposits and calcrete aquifers as mentioned for the Augustus subregion are of importance as these geologies are known to support subterranean habitats for both troglofauna and stygofauna (EPA 2016a).

Figure 3–1
Location of the Beyondie Potash Project in relation to IBRA regions and subregions



Client: Kalium Lakes Ltd
Project: Beyondie Sulphate of Potash Project
Author: AL
Date: 12-Jan-18
Coordinate System: GDA 1994 MGA Zone 51
Projection: Transverse Mercator
Datum: GDA 1994



3.2 CLIMATE AND WEATHER

The Gascoyne bioregion has an arid climate with and summer rainfall in the east. Spatially averaged median (1890–2005) rainfall is 202 mm (DEWHA 2008a). The climate of the Little Sandy Desert bioregion is also arid with summer-dominant rainfall. Spatially averaged median (1890–2005) rainfall is 178 mm (DEWHA 2008b). The climate of south-western Little Sandy Desert has also been described as desert tropical with predominant summer rainfall (van Leeuwen 2002).

The nearest Bureau of Meteorology (BoM) weather station with long-term and actual data is Newman Airport (BoM Station 7176; Latitude 23.42°S Longitude 119.80°E), approximately 160 km to the north-west of the study area. Newman records the highest maximum mean monthly temperature (39.1°C) in December and the lowest maximum mean annual temperature (22.8°C) in June. The lowest mean minimum temperature is recorded in July (6.4°C) and the highest in January (24.9°C). Average annual rainfall is 327.7.4 mm with February, January, and March recording the highest monthly averages (71.7, 67.5, and 44.0 mm respectively) (Figure 3-2).

Pan evaporation for the south-western Little Sandy Desert bioregion ranges from 16.1 mm/day in January to 4.5 mm/day in June at an annual daily average of 10.2 mm (van Leeuwen 2002).

The three months prior to the pilot survey recorded overall higher rainfall than average with a total of 359 mm from January to March 2017 providing ideal conditions for subterranean fauna surveys (Figure 3-2). Temperatures were about average in the year preceding the survey, although January 2017 was somewhat colder (Figure 3-2).




Figure 3-2 Average monthly temperatures and rainfall for Newman (BoM 2017)

3.3 GEOLOGY

The Project is located within the Collier, Salvation, Scorpion, and NW Officer Basins which reside within the Capricorn Orogeny, that marks the convergence and collision of the Archaean Pilbara and Yilgarn Craton and was responsible for widespread granite magmatism, deformation and metamorphism (Advisian 2017a).

3.3.1 Brine extraction area

Beyondie Lakes, Ten Mile Lake and Lake Sunshine are developed within a palaeochannel system (that includes regionally expansive salt lay playas) that contains the potassium rich sub-surface brines, hosted in Cenozoic colluvial deposits. The palaeochannel is underlain by bedrock of the Sunbeam Group, consisting mostly of sedimentary sandstones, siltstones, conglomerates and shales (Grey *et al.* 2005).

Intruded into the bedrock are dolerite intrusions (dykes and sills – Nd), for example expressed in the north-western part of the study area between the Beyondie freshwater marsh (just outside the study area) and eastern and western salt pan (Figure 3-3). Palaeochannels have been incised into the bedrock. Within the lakes, Quaternary lacustrine deposits (Qt) are mainly clay, mud and silt which are usually saline and commonly gypsiferous. Surrounding the lakes are expanses of Quaternary Aeolian sand and sand sheets (Qd) (Figure 3-3). Longitudinal (seif), chain and net dunes are abundant and there are some areas of ironstone pebble veneer.

The most prospective geologies in brine extraction areas for subterranean fauna are those of Quaternary alluvials (Qa) and calcretes (Czk) as identified in Figure 3-3.


Two main calcretes (Czk) are of potential relevance for the Project; these are present west and south of Ten Mile Lake consistent with two palaeochannels terminating at the lake system and currently reflected in two surface drainage lines, i.e. Nanyeriny Creek flowing into the Beyondie Lakes from the west and 477 Creek flowing into Ten Mile Lake from the south. All other calcrete bodies are located more than 5 km from the lakes and unlikely to be affected by brine extraction. One Quaternary alluvial (Qa) deposit is located adjacent to the one of the main calcretes (Figure 3-3).

3.3.2 Freshwater development envelope


The Freshwater Development Envelope encompasses a range of geologies but in general is targeting freshwater aquifers hosted in Quaternary surficial alluvial deposits. The alluvial deposits are generally formed as transported sediments during rain periods forming creek sediments and outwash fans on the flanks of the topographic lows and valleys. Thicker deposits of colluvium may also occur within tributaries in general break of slope areas and along steeper valley sides. These deposits are heterogeneous due to the nature of their deposition. Minor, localised calcretes are present as carbonate deposits formed within alluvial and colluvial sediments (Advisian 2017b).

Figure 3–3
Surface geology of the
Beyondie Potash Project

- Freshwater Development Envelope
- Tenement E69/3309
- Tenement E69/3346
- Tenement E69/3347
- Tenement E69/3351
- Tenement E69/3352
- Road
- Minor road
- Major creeks and rivers
- Lake



Client: Kalium Lakes Ltd
Project: Beyondie Sulphate of Potash Project
Author: AL
Date: 12-Jan-18
Coordinate System: GDA 1994 MGA Zone 51
Projection: Transverse Mercator
Datum: GDA 1994



3.4 HYDROLOGY

The four main aquifers that exist in the area are (Advisian 2017b; AQ2 2016):

- basal sands in the thalweg of the palaeochannel (target of the potassium brine resource)
- upper unconsolidated sediments in the palaeochannels (<10 m deep); i.e. alluvial (heterogenous, coarse) (targeted by the Freshwater Development Envelope) and lacustrine (gypsiferous, sand, silts, clays) deposits
- areas of saturated calcrete (especially to the west and south of Ten Mile Lake)
- areas of weathered or fractured bedrock.

3.4.1 Brine extraction area

Ten Mile Lake is a large salt playa and forms the western end of a chain of ephemeral salt lakes which extend eastwards and include Lake Sunshine, Yanneri Lake and Terminal Lake. This suite of lakes do not hydrologically connect above ground but they form part of the Ilgarari palaeoriver which is a remnant of an extensive river system from the tertiary period (Gentilli 1979).

The Ilgarari palaeoriver is a tributary of the Disappointment palaeoriver, that includes Lake Disappointment (Beard 2005). The catchment for Lake Disappointment is largely the Little Sandy Desert bioregion (Beard 2005). These salt lakes are dry most of the year but can become seasonally inundated during the wet season if there is sufficient rainfall and it is considered likely that some surface water recharges the underlying palaeochannel aquifers (AQ2 2016).

AQ2 (2016) note that the relationship between Beyondie and Ten Mile Lakes and the palaeochannel is important as it will influence the nature of brine abstraction from the palaeochannel system. For Beyondie and Ten Mile Lakes, it is considered likely that flow down the palaeochannel is controlled (on a local playa scale) by evaporative discharge (AQ2 2016). Deflation of exposed lakebeds along palaeochannels results in the lowering of the topographic elevation of lakebeds, thereby effectively bringing the groundwater level closer to the surface, promoting evaporation (AQ2 2016). The evaporative “pumping” increases groundwater discharge at the lake site, thus promoting groundwater flow towards the playa lakes.

Pumping of water from the basal sand aquifer may induce changes in the hydrostatic-heads within the different aquifer horizons and potentially induce flow from adjacent, i.e. sedimentary and calcrete aquifers (AQ2 2016).

3.4.2 Freshwater development envelope

As described in section 3.3.2, abstraction from the Freshwater Development Envelope will likely take place within aquifers containing alluvium and calcrete, which are heterogeneous due to the nature of their deposition (Advisian 2017b). Here, localised groundwater supplies may be obtained from intersections of present-day drainage, bedrock fracture zones and calcrete formation.

Where fractured bedrock supplies water to the alluvium, they can be open or closed and yields can be low to moderate and highly variable. In the unconfined alluvial zones groundwater will be rainfall dependent and will be higher in areas of highly heterogeneous coarser sediments. In windblown arid areas, the shallow alluvial sediments may have been eroded away to expose the calcrete deposits which have become hard outcrop features at the surface (Advisian 2017b).

Calcretes form an important water resource elsewhere, such as in the Goldfields region, where they can produce significant bore yields of up to 2,000 kilolitres per day (kL/d), if there is sufficient thickness

below the water table (Johnson *et al.* 1999). Calcrete is often associated with lower salinity water than other surrounding and deeper aquifers due to their ability to accept recharge. Accordingly, they are also significant subterranean fauna habitat in regions within the Yilgarn Craton (see section 3.5.4).

Calcrete occurs in the region at the margins of the present-day salt lakes, and in some of the main sub-catchments in the palaeodrainages. In the Project area the surficial unconfined aquifer comprising alluvium and colluvium associated within the present-day drainage, contains calcrete logged at maximum 6.57 m saturated thickness (south of Ten Mile Lake) and 5.27 m saturated thickness west of Beyondie Lakes (Advisean data in prep.).

3.5 BIOLOGICAL CONTEXT

Subterranean fauna lives within air- or water-filled underground networks. They are predominantly invertebrates. Organisms specialised for living in air-filled subterranean networks are referred to as troglotauna, while those inhabiting water-filled subterranean networks are referred to as stygofauna (Howarth 1983; Humphreys 2000).

Subterranean habitats are perpetually dark, are extremely constant in temperature and humidity (air-filled networks) and very low in nutrients and energy (Howarth 1993). Evolution under such conditions has resulted in much specialised organisms that are restricted to the void networks (Harvey 2002; Holsinger 2000; Howarth 1993; Ponder & Colgan 2002). Such species are obligated to living in subterranean networks and cannot live in epigean (surface) environments.

Organisms specialised to live in subterranean networks are likely to represent short-range endemics (SREs) with extremely limited capabilities of dispersal (Harvey 2002; Ponder & Colgan 2002; Volschenk & Prendini 2008). They are species with naturally small distributions, nominally less than 10,000 km², although species restricted to subterranean void systems may have considerably smaller distributions and therefore represent extreme SREs (Harvey 2002). It is these subterranean species that are considered to be of conservation significance because they are at greatest risk of extinction from project developments.

In WA, and particularly in the Pilbara and Midwest region, there has been a recent renaissance in the study of subterranean biodiversity (Humphreys 2008) driven by the growth of the mineral resources industry and mining environmental impact assessment (EIA) (EPA 2016a, c). However, the biology, diversity and distributions of most of WA's subterranean fauna are still poorly understood.

3.5.1 Stygofauna

Stygofauna represent the fauna living within subterranean water bodies or aquifers (Humphreys 2000). They typically show similar traits to troglobites in their specialisation to subterranean life, including loss of body pigment, eyes and heightened mechano-sensory systems. Stygofauna can be distinguished by their propensity for subterranean life (Humphreys 2008):

- stygobites — restricted to subterranean habitats and usually perish on exposure to the surface environment
- stygophiles — facultatively use subterranean habitats but are not reliant on them for survival
- stygoxenes — inhabitant of surface water which may also be able to freely move from surface to subterranean systems and back.

Short-range endemic stygofauna are only represented by stygobitic species.

3.5.2 Troglafauna

Troglafauna are typically divided into three categories of specialisation to subterranean life (Barr 1968; Howarth 1983; Humphreys 2000):

- troglobites — restricted to subterranean habitats and usually perish on exposure to the surface environment
- troglophiles — facultatively use subterranean habitats but are not reliant on them for survival
- troglonexes — use subterranean systems for specific purposes, such as roosts for reproduction (e.g. bats and swiftlets).

Both troglobites and troglophiles may be SREs and are therefore conservation significant.

Stygobites and troglobites are often characterised by specialised adaptations to subterranean life that allows them to exploit the dark, often nutrient-poor subterranean void networks, such as (Howarth 1983, 1993; Humphreys 2000; Poulson & Lavoie 2000):

- lack or reduction of eyes
- elongated limbs, but lack or reduction of wings (for species that are normally winged)
- lack or reduction of body pigmentation
- heightened chemosensory and mechano-sensory systems
- loss of circadian rhythms
- very low metabolic rate.

3.5.3 Threatening processes

Impacts to subterranean fauna can be classed as either:

- **primary impacts** – impacts that physically terminate the subterranean void networks
- **secondary impacts** – impacts that change the subterranean habitat without physically destroying the void networks.

Primary impacts are obvious, whereas secondary impacts tend to be cumulative and may affect a far greater area than that being developed (Hamilton-Smith & Eberhard 2000). There are commonly two key threatening processes from mining/extractive activities that impact subterranean fauna through the direct loss of habitat:

- **Development of mine pits** – the most obvious primary impact to subterranean habitats occurs as a result of their physical removal during mining. Troglafauna require air-filled void networks and most of this habitat exists in the overburden, which is typically destroyed during pit construction/excavation. Similarly, direct loss of stygofauna habitat may be caused by the removal of geological formations if any aquifers are associated with these formations. This Project is essentially a borefield, no direct removal of habitat will take place and therefore, this primary impact is not relevant.
- **Depletion of an aquifer leading to loss of stygofauna habitat** – depletion of an aquifer that is identified as suitable for stygofauna represents a direct loss of stygofauna habitat. The significance of the impact is dependent on the depth of drawdown, the size and extent of the aquifer and the connectivity of the aquifer with adjacent habitat for stygofauna.

Secondary impacts are those that affect the physicochemical properties of subterranean habitats. The nature of these changes can be difficult to measure and there is limited empirical evidence to support or refute these putative impacts. There are four secondary impacts that may be relevant to the Project:

- **Depletion of an aquifer leading to altered relative humidity** – troglofauna are dependent on high relative humidity (Barr 1968; Humphreys 1991; Humphreys 2000). Dewatering may impact troglofauna habitat in unsaturated strata above the water table by lowering relative humidity.
- **Nutrient starvation** – surface vegetation is the primary source of nutrients entering subterranean systems. Large-scale clearing of vegetation may result in the localised nutrient starvation of underlying subterranean habitat. Smothering of these nutrient sources on which subterranean systems depend, in the form of waste and overburden stockpiles and tailings ponds, may reduce inflow of nutrients to subterranean systems and lead to nutrient deficient habitats (Howarth 1993; Humphreys 2000; Poulson & Lavoie 2000).
- **Vibration** – shock waves through subterranean strata from blasting or heavy vehicle traffic may result in the collapse of less-consolidated void spaces and also impact physically on subterranean fauna. There is little data to challenge or corroborate these observations and impacts may generally be localised rather than critically threatening.
- **Contamination:** contamination of subterranean habitats from spills, such as diesel fuel, may degrade the quality of subterranean habitats. Such impacts would generally be highly localised and minor in scale; however, major contamination of subterranean habitats may have significant impacts.

3.5.4 Calcretes as subterranean fauna habitat

In WA, the types of geology known to support stygofauna include calcretes, alluvial formations particularly when associated with alluvial or palaeochannel aquifers, fractured rock aquifers and karst limestone. Troglofauna are likely to be present in karst, channel iron deposits (CID), banded iron formations (BIF), alluvium/colluviums in valley-fill areas, and weathered or fractured sandstone (EPA 2016a). Of these, alluvial or palaeochannel aquifers and to a lesser extent, calcretes are most prominent within the study area (Figure 3-3).

Calcretes are carbonate deposits that formed near the water table in arid lands as a result of concentration processes by near-surface evaporation. The calcretes are mainly associated with the palaeodrainage channels of the Pilbara and Yilgarn cratons and their associated orogens which together form the Western Shield of Australia (Beard 1998).

The study area is located immediately north (0-25 km) of the northern extent of the Yilgarn Craton and therefore strictly speaking does not occur within the craton; given the proximity however, the role of calcretes with respect to subterranean fauna are discussed in detail.

Calcrete deposits of the Yilgarn region of central WA are believed to have formed from the groundwater flow between 30–10 Mya when continued aridity produced salt lakes, extensive alluvial fans and dunes which assisted in fragmenting the river valleys into isolated ponds (Morgan 1993). In the northern river valleys of the Western Shield calcretes are concentrated on the upstream side of the salt lakes (Morgan 1993).

Over 200 major calcretes are found in the Yilgarn and they provide habitat for a diversity of subterranean fauna (both stygo- and troglofauna), such as diving beetles (Dytiscidae) and a variety of crustaceans, including amphipods (Amphipoda), slaters (Isopoda), syncarids (Bathynellacea) and copepods (Copepoda) (Bradford *et al.* 2013; Cooper *et al.* 2008; Guzik *et al.* 2008; Humphreys 2001;

Javidkar *et al.* 2015; Javidkar *et al.* 2016; Javidkar *et al.* 2017; Watts & Humphreys 2004). The calcretes are generally shallow (approximately up to 10 m deep), but may reach depths of up to 30 m (Humphreys *et al.* 2009).

The calcretes of the Western Shield that occur near salt lakes have recently been flagged as 'groundwater estuaries' of potentially high biodiversity value (Humphreys *et al.* 2009). Their groundwater profiles can be compared to the anchialine systems near ocean estuaries, where saline water intrudes under the freshwater fan of the river mouth. The complex interplay of fresh and saline water creates steep biogeochemical gradients to potentially support cascades of microbiological and micro- and macroinvertebrate communities (Humphreys *et al.* 2009).

The high endemism of the groundwater stygofauna of calcrete deposits in the Yilgarn Craton has resulted in the listing of 77 PECs for the Midwest and the Goldfields regions of WA (DPaW 2016b) (Appendix 3).

3.5.5 Salinity

Globally, stygofauna are generally restricted to freshwater and rarely occur in mildly brackish waters, except in the special case of anchialine ecosystems (Bradbury & Williams 1996b; Humphreys 1999). However, the Western Shield contains a diverse assemblage of near-marine and ancient freshwater lineages inhabiting groundwater with salinity values that may reach marine conditions (Humphreys 2008). At Lake Way, Watts and Humphreys (2004) recorded a diverse stygal assemblage in a bore with surface salinity near seawater (30,000 mg/L) and a strong salinity gradient increasing to 69,000 mg/L at 6 m depth. The finer scale vertical distribution of stygofauna inhabiting groundwater in WA, which may be strongly stratified with steep gradients in salinity at micro/meso-scales, remains largely unknown (Humphreys *et al.* 2009; Subterranean Ecology 2010b).

4 METHODS

4.1 DESKTOP REVIEW

This assessment followed EPA guidelines which state that desktop reviews for subterranean fauna should (EPA 2016a):

- search regional and project/site specific habitat data, including geological and hydrological information, previous studies of the area (published and unpublished), site photographs and databases including fauna records
- place the Project area into a regional context
- make conclusions about whether the area is likely to provide habitat for subterranean fauna and consider impacts of the proposal.

4.1.1 Database searches

Database searches and literature reviews of relevant publications were undertaken to compile a list of conservation significant subterranean species that may occur within the study area based on the proximity of previous records.

The following database searches were undertaken for a quadrat of approximate 100 km length with the diagonal coordinates of -24.31222°S, 119.78444°E (NW point) and -25.23472°S, 120.7808°E (SE point):

- EPBC Act Protected Matters Search Tool (Department of the Environment 2015)
- DPaW/WA Museum NatureMap (DPaW 2015a)
- DPaW Threatened Flora, Fauna and Ecological Communities databases (DPaW 2015c).

In addition, the WA Museum Arachnology/Myriapodology, Crustacea and Mollusca databases were undertaken for a quadrat of approximate 200 km length, consistent with the nominal range of short-range endemic invertebrates (EPA 2016d), with the diagonal coordinates of -23.86°S, 119.30°E (NW point) and -25.67°S, 121.27°E (SE point).

4.1.2 Literature review

Several subterranean fauna surveys or desktop reviews near salt lakes were consulted to provide context of biodiversity for large inland salt lakes in WA, including Lake Disappointment, Lake Way and Lake Lefroy (Table 4-1). It is recognised that these lakes are a considerable distance from the study area; however, any salt lake in the state can provide background data to contextualise the desktop review and subsequent survey results. All of these salt lakes are also under some impact from mining activities.

Table 4-1 Selected aquatic invertebrate surveys of salt lakes in Western Australia

Project	Author	Remarks
Lake Way	Toro Energy (2011, 2012)	Closest to study area and similar geological setting (calcrete near salt lakes) but located within the Yilgarn Craton; very comprehensive survey program over a number of years.
Lake Disappointment	Phoenix (2014)	Subterranean desktop assessment only; located in same palaeochannel as the study area.
Lake Lefroy	Subterranean Ecology (2010a, b)	Separate troglotauna and stygofauna assessments; located within the Yilgarn Craton.

4.2 FIELD SURVEY

4.2.1 Survey


A total of 15 bores were surveyed for subterranean fauna including seven within the exploration tenements and eight regional reference sites (Table 4-2; Figure 4-1) between 29–31 March 2017.

Table 4-2 Location and details of bores sampled during field survey


Bore	Latitude (GDA94)	Longitude (GDA94)	Depth (m)	Depth to water (m)
David's Well	-25.030062	120.007066	21	6
No.77 East Well	-24.975719	120.189032	10	1
Broken Leg Bore	-24.997021	120.056049	42	16
Garden Well	-24.782386	120.061737	22	11
Beyondie Bore	-24.782565	120.043510	24	11
Tupree Well	-24.660603	120.054702	13	9
12 Mile Well	-24.731820	120.224108	11	5
Unnamed Bore	-24.769961	120.111655	20	4
WM05MBS	-24.732438	120.326538	53	3
WB14 (= Beyondie West)	-24.784564	120.036625	>100	11
WB25	-24.767690	120.384670	29	7
WB09MB5	-24.792840	120.333731	31	3
TMAC23	-24.809518	120.338262	12	3
WB12MB1	-24.796494	120.367315	34	1
WB10TB1	-24.766475	120.363892	74	7

Figure 4-1
Location of bores
sampled during
field survey

- Freshwater Development Envelope
- Tenement E69/3309
- Tenement E69/3346
- Tenement E69/3347
- Tenement E69/3351
- Tenement E69/3352
- Road
- Minor road
- Major creeks and rivers
- Lake
- Bores



Client: Kalium Lakes Ltd
Project: Beyondie Sulphate of Potash Project
Author: AL
Date: 12-Jan-18
Coordinate System: GDA 1994 MGA Zone 51
Projection: Transverse Mercator
Datum: GDA 1994



Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community

4.2.2 Fauna sampling

The subterranean fauna survey was conducted using six stygofauna hauls, three with a 250 µm and three with a 50 µm weighted nets (diameter 90 mm). After the net was lowered to the bottom of each bore, it was used to briefly stir up sediments and their benthic inhabitants. All nets were equipped with nylon 'ticklers' to collect any troglofauna animals moving along the sides of the bores. The nets were scraped back up the bore wall with each scrape covering a different part of the bore wall.

After each haul/scrape, the strained content was rinsed into a 120 ml plastic vial by squirting 100% ethanol down the sides of the net and around the rim of the weight, washing the sample contents into the vial. If not already full, the sample vial was topped up with 100% ethanol.

The net was thoroughly rinsed in freshwater after each sample to avoid cross-contamination of samples.

At conclusion of the survey, samples were stored in a refrigerator in the laboratory, where they were sorted and specimens identified using high-magnification stereo-microscopes.

4.2.3 Bore and water quality data

At each bore, depth and depth to groundwater were measured (see Table 4-2). Water quality parameters were measured in-situ with a YSP multiprobe in a water sample bailed up from the bore and included:

- temperature (°C)
- dissolved Oxygen (%)
- dissolved Oxygen (mg/L)
- conductivity (µs/cm)
- salinity (ppt)
- pH
- oxygen reduction potential (mV).

4.2.4 Morphological species identifications

Phoenix has considerable in-house expertise in the identification of SRE target groups. Senior staff involved in identifications are also Research Associates with a longstanding taxonomic research history at the WA Museum. For some groups, external experts were consulted (Table 4-3).

4.2.5 Molecular identifications

The identification of species based on comparisons between DNA sequences is referred to as DNA barcoding and is an expected identification tool in survey of subterranean fauna (EPA 2016a). Any gene can be used for barcoding purposes; however, the primary gene targeted by researchers is Cytochrome Oxidase Subunit I (COI) (Hebert *et al.* 2003).

Molecular identifications were conducted by staff of Helix Molecular Solutions, based at the School of Biological Sciences, University of Western Australia (Table 4-3).

4.3 PROJECT PERSONNEL

The personnel involved in the survey are presented (Table 4-3).

Table 4-3 Project team

Name	Qualifications	Role/s
Dr Volker Framenau	B.Eng. (Chem. Eng.), M.Sc. (Cons. Biol.), Ph.D. (Zool.)	Project manager, taxonomy (Araneae, Hemiptera), report writing
Mr Jarrad Clark	B.Sc. (Env. Sci.)	Field survey, report writing
Ms Karen Crews	B.Sc. (Env. Sci.) (Hons)	Report review
Mrs Kathryn Wyatt	BIS. (GIS), Grad. Cert. (GIS)	GIS
Ms Jane McRae ¹		Taxonomy (Copepoda, Cladocera, Conchostraca)
Ms Yvette Hitchen ²		Molecular analyses (laboratory)
Dr Terrie Finston ²		Molecular analyses (analyses, report writing)

1 – Bennelongia; 2 – Helix Molecular Solutions.

5 RESULTS

5.1 DESKTOP REVIEW

5.1.1 Database searches

The database searches did not return any subterranean records from the study area, but returned seven species of stygofauna in three orders of crustaceans (Bathynellacea, Cyclopoidea, Podocopida) from within approximately 100 km of the Project (Table 5-1; Figure 5-1). Three of these species are currently known from single localities and the remaining four species are known to be widespread (see section 5.2).

No troglofauna records were returned in the database searches.

5.1.2 Threatened and Priority species or ecological communities

No Western Australian subterranean fauna or ecological communities are listed as matters of NES (Department of the Environment 2014a).

No Threatened or Priority subterranean fauna species or ecological communities are listed from the Little Sandy Desert (DPaW 2016a, b, 2017) (see also Appendices 1–3). Many subterranean stygofauna communities are listed as PECs from calcretes of the Yilgarn Craton; however, the study area does not occur within the Yilgarn Craton.

5.1.3 Literature review

Stygofauna and troglofauna were recorded for the Wiluna Uranium Project situated at Lake Way, a large inland salt lake in the Murchison region of WA approximately 230 km south-west of the study area (Toro Energy 2011, 2012). At least 67 stygofauna but only up to 20 troglofauna species were recorded from the surveys encompassing the Hinkler Well, Lake Violet and Uramurdah calcretes groundwater systems, all listed as PECs (DPaW 2015b). The salinity levels in the upper strata of the bores where some of these species were recorded were up to 141,200 $\mu\text{S}/\text{cm}$ (hypersaline). Two key factors relevant to the current Project were noted:

1. Several species recorded were found in more than one calcrete system, challenging the notion that calcretes are isolated subterranean islands.
2. No subterranean samples were collected from the playa of Lake Way.




A subterranean desktop study for a proposed Potash mine at Lake Disappointment (280 km north-east of the study area) concluded that there was low likelihood of subterranean fauna to be present under the lake playa due to high salinity levels and low porosity lacustrine sediments (Phoenix 2014). However, calcrete formations to the west of Lake Disappointment were considered as subterranean fauna habitat and surveys were recommended if these calcretes were likely to be affected by the development.

Detailed desktop studies for subterranean fauna at Lake Lefroy identified variable likelihoods of stygo- and troglofauna to occur based on geology (porosity/permeability of the sediment) and, for stygofauna, salinity (Subterranean Ecology 2010a, b). Overall, the likelihood of stygofauna (Subterranean Ecology 2010a) or troglofauna (Subterranean Ecology 2010a) to occur was low; however, the Goldfields region lacks the calcrete deposits typical for the Yilgarn.



Table 5-1 Subterranean fauna recorded in the desktop review


Family	Genus and species	Troglofauna/ stygo fauna	SRE rating	Locality	Latitude (GDA94)	Longitude (GDA94)
Syncarida – Bathynellacea						
Parabathynellidae	<i>Billibathynella ilgarariensis</i>	Stygo fauna	SRE	Bulloo Downs Station, Ilgarari Creek, Yanneri Well	-24.439444	119.7575
	<i>Brevisomabathynella magna</i>	Stygo fauna	SRE	Cunyu Station, Sweetwaters Well, Nabberu Palaeodrainage	-25.610556	120.3725
Copepoda – Cyclopoidea						
Cyclopidae	<i>Pilbaracyclops frustratio</i>	Stygo fauna	SRE	Yanneri Well, Bulloo Downs Station, Ilgarari Creek, Lake Disappointment	-24.439722	119.7575
	<i>Mesocyclops brooksi</i>	Stygo fauna/ stygo phile	Widespread	No. 37 Government Well, Bulloo Downs Station, Ilgarari Creek	-24.313611	119.6975
				Well (Site 31), Bulloo Downs Station, Ilgarari Creek	-24.327778	119.672778
				Old Ilgarari Well, Bulloo Downs Station, Ilgarari Creek	-24.334167	119.68
	<i>Microcyclops varicans</i>	Stygo fauna	Widespread	No. 37 Government Well, Bulloo Downs Station, Ilgarari Creek	24.313611	119.6975
Ostracoda – Podocopida						
Candonidae	<i>Deminutiocandona quasimica</i>	Stygo fauna	Widespread	Bulloo Well, Bds001, Pilbara region	-24.008333	119.575
Cyprididae	<i>Cyprinotus cingalensis</i>	Stygo fauna/ surface	Widespread	Government Well #40, Savory 2, Pilbara region	-23.863611	120.150667

Figure 5-1
Subterranean fauna
records from the
desktop review


-  Freshwater
-  Tenement E69/3309
-  Tenement E69/3346
-  Tenement E69/3347
-  Tenement E69/3351
-  Tenement E69/3352
-  Road
-  Minor road
-  Major creeks and rivers
-  Lake

Desktop records

-  SRE
-  Widespread



Client: Kalium Lakes Ltd
 Project: Beyondie Sulphate of Potash Project
 Author: AL
 Date: 12-Jan-18
 Coordinate System: GDA 1994 MGA Zone 51
 Projection: Transverse Mercator
 Datum: GDA 1994



5.2 FIELD SURVEY

5.2.1 Water quality

Water quality parameters varied greatly between bores, in particular with respect to salinity Table 5-2). Five bores had freshwater, four were subsaline, one hyposaline, one mesosaline and four contained hypersaline water. Water in most bores was slightly alkaline, with the exception of Broken Leg Bore and WM05MBS, which was both slightly acidic Table 5-2).

Table 5-2 Water quality data for sampled bores

Bore	Temperature (°C)	Dissolved Oxygen (%)	Dissolved Oxygen (mg/L)	Conductivity (µs/cm)	Salinity (ppt)	pH	Oxygen Reduction Potential (mV)
David's Well	28.5	80.6	5.7	724	0.34	7.37	131
No. 77 East Well	24.3	10	0.74	149.9	0.07	8.32	113.2
Broken Leg Bore	27.3	5	0.34	1,016	0.48	6.97	-166
Garden Well	25.5	54	4.16	804	0.38	7.5	23.8
Beyondie Bore	26.1	34.7	2.52	577	0.27	7.87	47.4
Tupée Well	26.1	14.3	1.06	5,980	3.0	7.51	110
12 Mile Well	28.8	89.2	6.21	3,238	1.62	7.92	82
Unnamed Bore	28.2	24.7	1.79	3,257	1.63	7.49	89.7
WM05MBS	28.8	26.6	1.34	95,128	65.45	6.83	197
WB14 (= Beyondie West)	26.4	59.6	4.52	1,593	0.77	8.22	91.7
WB25	27.3	20.4	1.37	26,650	15.62	7.35	137.7
WB09MBS	26	6.4	0.31	96,105	66.25	7.26	170.7
TMAC23	27.1	32.8	2.3	11,061	6.01	7.84	103.9
WB12MB1	28.5	9.2	0.35	144,185	108.47	7.06	149.7
WB10TB1	27.6	15.8	0.51	150,500	114.51	7.71	134.8

1 – Salinity (Hammer 1986): <0.5 – freshwater; 0.5–3 ppt – subsaline; 3–20 ppt – hyposaline; 20–50 ppt – mesosaline; >50 ppt - hypersaline

5.2.2 Subterranean fauna

A total of 11 taxa of aquatic invertebrates were recorded from the field survey, some in considerable numbers (Table 5-3). Only two are considered stygobites, paramelitid amphipods from Garden Well and Tupee Well, and syncarid crustaceans (order Bathynellacea) from Garden Well. The copepod *Mesocyclops brooksi* has initially been described from groundwater, but has subsequently been found mainly in wells and is now considered a stygophile rather than a stygobite (Karanovic 2006). All other aquatic organisms are widespread and often found in surface lentic (still water) systems (Table 5-3).

Table 5-3 Potential stygofauna fauna collected during field survey

Higher order	Genus and species	No. of specimens	Bores	Ecotype
Amphipoda (amphipods)				
Paramelitidae	Paramelitidae sp. indet.	8	Garden Well, Tupee Well	Stygobitic
Syncarida (syncarids)				
Parabathynellidae	Parabathynellidae sp. indet.	6	Garden Well	Stygobitic
Cladocera (water fleas)				
Daphniidae	<i>Ceriodaphnia quadrangula</i> s. l.	>100	No 77 East Well	Surface
Macrothricidae	<i>Macrothrix capensis</i>	1	Beyondie Bore	Surface
Moinidae	<i>Moina micrura</i> s. l.	>100	No. 77 East Well	Surface
Conchostraca				
Leptestheriidae	<i>Leptestheria</i> 'B01'	6	No. 77 East Well	Surface
Copepoda				
Cyclopidae	<i>Mesocyclops brooksi</i>	>100	Beyondie Bore, No. 77 East Well	Stygophile ; widespread
	<i>Mesocyclops</i> sp. indet.	2	No. 77 East Well, Tupee Well	Unknown; immature specimens
	<i>Thermocyclops decipiens</i>	>100	Beyondie Bore, No. 77 East Well	Surface; widespread
Diptera (flies)				
Chironomidae	Chironomidae sp. indet.	1	Beyondie Bore	Surface; not further identified as irrelevant to study
Oligochaeta (worms)				
Oligochaeta fam. indet.	Oligochaeta sp. indet.	2	Broken Leg	Unknown


A total of 17 specimens of potential troglofauna were collected, but only an unidentified spider from Garden Well and a meenoplid bug from David's Well are here considered to be troglobitic (Table 5-4). The remainder, mites and springtails, potentially belong to the edaphic (soil) fauna and are not further considered here.

Table 5-4 Potential troglofauna fauna collected during field survey


Higher order	Genus and species	No. of specimens	Bores	Ecotype
Acari (mites)				
Acari fam. indet.	Acari sp. indet.	13	Garden Well, Broken Leg, W12MBI	Troglophiles/edaphic; not considered in subterranean studies
Araneae (spiders)				
Araneomorphae fam. indet.	Araneomorphae sp. indet.	1	Garden Well	Troglobitic ; poor condition, family identification not possible
Hemiptera (bugs)				
Meenoplidae	Meenoplidae sp. indet.	1	David's Well	Troglobitic ; immature larvae
Collembola (springtails)				
Collembola fam. indet.	Collembola sp. indet.	2	Beyondie Bore, WB09MDB	Troglophiles/edaphic; not considered in subterranean studies

Figure 5-2
Subterranean fauna
collected during
field survey

- Freshwater Development Envelope
- Tenement E69/3309
- Tenement E69/3346
- Tenement E69/3347
- Tenement E69/3351
- Tenement E69/3352
- Road
- Minor road
- Major creeks and rivers
- Lake
- Troglofauna records
- Stygofauna records



Client: Kalium Lakes Ltd
Project: Beyondie Sulphate of Potash Project
Author: AL
Date: 12-Jan-18
Coordinate System: GDA 1994 MGA Zone 51
Projection: Transverse Mercator
Datum: GDA 1994



5.2.2.1 Araneae (spiders)

Spiders include a considerable number of subterranean forms in a number of families, in Australia represented by, amongst others, Trochanteriidae (Platnick 2008), Oonopidae (Baehr *et al.* 2012), Tetrablemmidae (Burger *et al.* 2010), Symphytognathidae (Harvey 2001b), Gnaphosidae (Phoenix, unpublished data) and others (Harvey *et al.* 1993). Currently, all known species of troglobitic spiders have a very narrow distribution and are often known from single bores or caves.

A poorly preserved, likely troglobitic small juvenile spider was recovered from Garden Well and is here considered to represent a troglobitic form due to colouration and lack of eyes (Table 5-4). It mostly resembles a symphytognathid spider, but accurate identification even at family level is not possible.

5.2.2.2 Hemiptera (bugs)

A single specimen of immature Meenoplidae was collected at David's Well (Table 5-4). It was classified as troglofauna as it is eyeless and troglobitic forms in the family are known from WA (Hoch 1993). However, the life cycles of species in the family are poorly known and it is possible that eyeless larvae have surface dispersal stages (Bennelongia 2012).

5.2.2.3 Amphipoda (amphipods)

There is a considerable diversity of subterranean amphipods world-wide (Holsinger 1993). The Western Australian fauna includes members of a number of families, such as Melitidae, Neoniphargidae, Paramelitidae, Bogidiellidae and others (Bradbury 2002; Bradbury & Williams 1997). Calcrete aquifers of the Yilgarn host a highly diverse amphipod fauna within the Hyalidae and Paramelitidae (Cooper *et al.* 2007).

Stygobitic amphipods in the family Paramelitidae were collected from Garden Well and Tupee Well (Table 5-3). They were submitted for molecular analyses and three of four specimens were successfully sequenced, all from Garden Well. Apart from length differences, all three sequences were genetically identical and therefore represent the same species (AMP036) in the family Paramelitidae. Phylogenetic analysis of the sequenced specimens and 43 reference sequences showed the lineage of the sequenced specimens differed from reference specimens by >15% sequence divergence, indicating they represent a species that has not previously been recorded in WA.

As specimens from Tupee Well could not be sequenced, it is not clear if they represent the same species (AMP036) that was recorded from Garden Well.

5.2.2.4 Syncarida (syncarids)

Syncarid crustaceans are often confined to subterranean environments. They are considered rare and difficult to collect and therefore very little is known about the biology and relationships (Abrams *et al.* 2013). They are regularly collected in calcretes of the Yilgarn and Pilbara cratons (Cho & Humphreys 2010; Guzik *et al.* 2008).

Two parabathynellids were recorded in the desktop review, *Billibathynella ilgarariensis* and *Brevisobathynella magna* (Table 5-1; Figure 5-1). Both are currently only known from type localities, ca. 65 km north-west and approximately 95 km south of the study area (Figure 5-1) (Cho & Humphreys 2010; Hong & Cho 2009).

Six specimens of syncarids were collected from Garden Well. Three of four specimens submitted for molecular analysis were successfully sequenced. Apart from length differences, all three sequences were genetically identical and therefore represent the same species (BAP028) in the family

Parabathynellidae. Phylogenetic analysis of the sequenced specimens and 69 reference sequences showed the lineage of the sequenced specimens differed from reference specimens by >15% sequence divergence, indicating they represent a species that has not previously been recorded in WA.

5.2.2.5 Copepoda (copepods)

Copepods are microscopic teardrop-shaped crustaceans that are common in freshwater and saline wetlands (Maly *et al.* 1997; Stoch 2001). They also form a considerable element of subterranean stygofauna in WA (Karanovic 2006).

Three subterranean species of copepods were identified in the desktop review, including one species only known from a single locality, *Pilbaracyclops frustratio* from approximately 65 km north-west of the study area (Table 5-1; Figure 5-1).

Only one of the copepod species identified from the desktop review was collected during the field survey, *Mesocyclops brooksi*. This species was described from ground waters of the Cape Range karsts (Pesce *et al.* 1996), but records have been forthcoming from the south-west of WA and interstate (Karanovic 2006). The species is considered a stygophile rather than stygobite due to its common presence in open wells rather than bores (Karanovic 2006). The species were collected at Beyondie Bore and No. 77 East Well (Table 5-1; Figure 5-1).

5.3 SURVEY LIMITATIONS

The EPA *Technical Guide: Terrestrial fauna surveys* (EPA 2016e) identifies potential limitations that may be encountered during terrestrial fauna surveys. These were applied here as no equivalent guidance with respect to subterranean fauna exists (Table 5-5). With respect to this survey, limited contextual information from around the study area was identified as the main limitation.

Table 5-5 Survey limitations based on EPA Technical Guide: Terrestrial fauna surveys (EPA 2016e)

Limitations	Limitation for this survey?	Comments
Competency/experience of survey personnel, including taxonomy	No	The field and laboratory teams and report authors have extensive experience in survey of subterranean systems in WA. Taxonomically poorly known groups were identified by molecular methods.
Scope and completeness - were all planned survey methods implemented successfully, was the study area fully surveyed	No	Suitable collecting methods were used based on comparable surveys in WA and consistent with subterranean pilot surveys conducted in WA.
Intensity - in retrospect, was the intensity adequate	No	The survey intensity of 15 bores was appropriate for a pilot subterranean fauna survey within the study area.
Proportion of fauna identified, recorded and/or collected	No	Fauna specimens collected match geological and hydrological conditions (i.e. no stygofauna in hypersaline water). All fauna specimens were identified to lowest possible taxonomic level.
Availability of adequate contextual information	Yes	There is good regional contextual information in relation to subterranean surveys in the Goldfields (in particular with respect to the Yilgarn) and Pilbara regions. However, little information was available near the study area (Gascoyne and Little Sandy Desert bioregions) as evidenced by the results of the desktop review.
Timing, weather, season, cycle	No	The survey was conducted at a suitable time of the year and soon after above average rainfall.
Disturbances which affected the results of the survey	No	No disturbances occurring during the period of the field survey are considered to have impacted the results.
Remoteness and/or access problems	No	There were no access problems in the study area.

6 DISCUSSION

The EPA's objective for subterranean fauna is its protection so that biological diversity and ecological integrity are maintained (EPA 2016b). Subterranean communities are often restricted to very small areas and it is supposed this is based on the limited dispersal capabilities of the fauna, with short-range endemism interpreted at a much smaller scale than in terrestrial systems (Eberhard *et al.* 2009).

The study area is situated in a poorly surveyed area where the eastern Gascoyne and south-western Little Sandy Desert bioregions intercept and where limited contextual information exists (van Leeuwen 2002). This was evident in the poor return from the desktop review that only revealed records of seven species of stygofauna and no troglofauna records.

Several factors were considered in assessing the likelihood of subterranean fauna occurrence in the study area, in particular the geological and hydrological setting of the Project, if there were subterranean fauna previously recorded near the Project and the results of surveys in similar conditions (i.e. alluvial and calcrete deposits near salt lakes in palaeovalleys in WA).

The desktop review findings indicate a high likelihood of occurrence of subterranean fauna in the Freshwater Development Envelope (Gascoyne bioregion) component of the study area, where freshwater aquifers in unconfined alluvial deposits are to be targeted.

The likelihood of subterranean fauna being present within the brine extraction areas is less clear from the desktop review findings. On one hand the study area is located on the border of the Yilgarn Craton, which has recorded highly diverse and restricted subterranean communities (e.g. Cooper *et al.* 2008; Watts & Humphreys 2001), and displays similar geological conditions, with the formation of alluvial deposits and calcrete expressions in ancient palaeochannels. On the other hand, the local geology (low porosity lacustrine deposits of fine mud, silt and clay) and hydrology (hypersaline) suggested a low likelihood of occurrence.

The pilot survey sampled bores and wells from both the Freshwater Development Envelope and brine extraction areas and the results largely affirmed the findings of the desktop review. Troglofauna and stygofauna were recovered from the alluvial deposits of the Nanyeriny Creek system west of Beyondie Lakes at Garden Well (freshwater), and Tupee Well (subsaline) and in a south-western reference bore, David's Well (freshwater), all within the Freshwater Development Envelope. Stygofauna records from the desktop review were also from alluvial deposits within the Gascoyne bioregion, approximately 65 km north-west of the study area. No subterranean fauna was recorded from bores associated with the brine extraction areas.

Molecular analysis confirmed the amphipods and syncarids from Garden Well are not conspecific with any previously sequenced species from WA and therefore represent new species. Only one species, the widespread stygophile copepod *Mesocyclops brooksi* was present in both the desktop review and field survey. It is noted that the phylogenetic analysis was limited by the lack of reference material from the region.

It is possible that the Garden Well and Tupee Well occur within a connected alluvial aquifer, which would be supported if the amphipods found at both sites represent the same species. As the Tupee Well specimen could not be sequenced, additional sampling/specimens would be required to confirm this.

Sampling associated with Ten Mile Lake (both in depositional sands and calcrete) failed to record subterranean fauna. As expected the low porosity lacustrine deposits (clay, mud and silt) associated with this lake and salinities in the brine in excess of 70,000 mg/L TDS, with levels consistently around 200,000 mg/L and sometimes in excess of 300,000 mg/L (up to 114.1 ppt in the sampled bores; see Table 5-2) have likely prohibited the presence of subterranean fauna (particularly stygofauna).

It is noted however that sampling in the calcrete to the south of Ten Mile Lake (considered the most prospective subterranean habitat in that area prior to sampling), was limited to a single bore (TMAC23). Here the salinity was subsaline and therefore suitable for stygofauna to occur, but porosity was extremely low, with fine clays recovered during sampling causing the water to take over one hour to drain from the sample sieves. This calcrete formation is very large, being approximately 15 km in length (north-south) and up to 7.5 km in width. The maximum saturated thickness is a relatively thin 6.57 m, at approximately 7.5 km SW of Ten Mile Lake (Advisian data in prep.). Current modelling based on pump tests suggests that after 23 years of brine extraction at Ten Mile Lake, groundwater will be lowered by 1– 2 m, at a distance 1-2 km across the entirety of this calcrete.

6.1 RECOMMENDATIONS

The survey confirmed the presence of subterranean fauna within the alluvial areas being targeted for the supply of freshwater for processing. Very little is known about the subterranean fauna species composition, distribution and degree of endemism in the region. It is recommended that the survey effort is increased to level 2 for stygofauna consistent with EPA (2016a), once a sufficient number of bores are available for sampling.

The survey need not survey troglofauna as no significant impacts to troglofauna are expected under the current plans. The expected limited drawdown of aquifers associated with the Freshwater Development Envelope are unlikely to significantly affect troglofauna. In addition, no troglofauna are expected to occur in association with Ten Mile Lake or Lake Sunshine, therefore the brine extraction process will not impact troglofauna.

7 REFERENCES

- Abrams, K. M., King, R. A., Guzik, M. T., Cooper, S. J. B. & Austin, A. D. 2013. Molecular phylogenetic, morphological and biogeographic evidence for a new genus of parabathynellid crustaceans (Syncarida : Bathynellacea) from groundwater in an ancient southern Australian landscape. *Invertebrate Systematics* **27**: 146–172.
- Advisian. 2017a. *Beyondie Potash Project - Ten Mile and Sunshine Lakes. Hydrological assessment of brine abstraction*. Advisian, West Perth, WA. Unpublished report prepared for Kalium Lakes Pty Ltd
- Advisian. 2017b. *Beyondie Potash Project. Preliminary water supply assessment*. Advisian, WorleyParsons Group, West Perth, WA. Unpublished report prepared for Kalium Lakes Potash Pty Ltd.
- AQ2. 2016. *Beyondie Lakes Project. H2 Level of assessment*. AQ2 Pty Ltd, Perth, WA. Unpublished report prepared for Kalium Lakes Potash Pty Ltd.
- Baehr, B., Harvey, M. S., Burger, M. & Thoma, M. 2012. The new Australasian goblin spider genus *Prethopalpus* (Araneae, Oonopidae). *Bulletin of the American Museum of Natural History* **369**: 1–113.
- Barr, T. C. 1968. Cave ecology and the evolution of troglobites. In: Dobzhansky, T., Hecht, M. K. & Steere, W. C. (eds) *Evolutionary biology*. North-Holland, Amsterdam, pp. 35–102.
- Beard, J. S. 1998. Position and developmental history of the central watershed of the Western Shield, Western Australia. *Journal of the Royal Society of Western Australia* **81**: 157–164.
- Beard, J. S. 2005. Drainage evolution in the Lake Disappointment catchment, Western Australia - a discussion. *Journal of the Royal Society of Western Australia* **88**: 57–64.
- Bennelongia. 2012. *Addendum: Troglofauna assessment, Ularing haematite Project*. Bennelongia Pty Ltd, Jolimont, WA. Unpublished report prepared for Macarthur Minerals Ltd.
- BoM. 2017. *Climate statistics for Australian locations*. Commonwealth of Australia, Bureau of Meteorology. Available at: <http://www.bom.gov.au/climate/data/>
- Bradbury, J. H. 2002. Melitid amphipods of Barrow Island, Western Australia. Part II - recent discoveries. *Records of the Western Australian Museum* **21**: 83–103.
- Bradbury, J. H. & Williams, W. D. 1996a. Freshwater amphipods from Barrow Island, Western Australia. *Records of the Australian Museum* **48**: 33–74.
- Bradbury, J. H. & Williams, W. D. 1996b. Two new species of anchialine amphipod (Crustacea: Hadziidae: *Liagoceradocus*) from Western Australia. *Records of the Western Australian Museum* **17**: 395–409.
- Bradbury, J. H. & Williams, W. D. 1997. The amphipod (Crustacea) stygofauna of Australia: description of new taxa (Melitidae, Neoniphargidae, Paramelitidae) and a synopsis of known species. *Records of the Australian Museum* **49**: 249–341.
- Bradford, T. M., Adams, M., Guzik, M. T., Humphreys, W. F., Austin, A. D. & Cooper, S. J. B. 2013. Patterns of population genetic variation in sympatric chiltoniid amphipods within a calcrete aquifer reveal a dynamic subterranean environment. doi:10.1038/hdy.2013.22.
- Burger, M., Harvey, M. S. & Stevens, N. 2010. A new species of blind subterranean *Tetrablemma* (Araneae: Tetrablemmidae) from Australia. *The Journal of Arachnology* **38**: 146–149.
- Car, C. A., Short, M., Huynh, C. & Harvey, M. S. 2013. The millipedes of Barrow Island, Western Australia (Diplopoda). *Records of the Western Australian Museum, Supplement* **83**: 209–219.
- Cho, J. L. & Humphreys, W. F. 2010. Ten new species of the genus *Brevisomabathynella* Cho, Park and Ranga Reddy, 2006 (Malacostraca, Bathynellacea, Parabathynellidae) from Western Australia. *Journal of Natural History* **44**: 993–1079.
- Cooper, S. J. B., J.H. Bradbury, Saint, K. M., Leys, R., Austin, A. D. & Humphreys, W. F. 2007. Subterranean archipelago in the Australian arid zone: mitochondrial DNA phylogeography of amphipods from central Western Australia. *Molecular Ecology* **16**: 1533–1544.

- Cooper, S. J. B., Saint, K. M., Taiti, S., Austin, A. D. & Humphreys, W. F. 2008. Subterranean archipelago: mitochondrial DNA phylogeography of stygobitic isopods (Oniscidea: *Haloniscus*) from the Yilgarn region of Western Australia. *Invertebrate Systematics* **22**: 195–203.
- Cowan, M. & Kendrick, P. 2001. Little Sandy Desert 2 (LSD2—Trainor subregion). In: May, J. E. & McKenzie, N. L. (eds) *A biodiversity audit of Western Australia's 53 biogeographical subregions in 2002*. Department of Conservation and Land Management, Perth, WA, pp. 413–422.
- Danielopol, D. L., Baltanás, A. & Humphreys, W. F. 2000. *Danielopolina kornickeri* sp. n. (Ostracoda, Thaumatocypridoidea) from a western Australian anchialine cave: morphology and evolution. *Zoologica Scripta* **29**: 1–16.
- Department of the Environment. 2014a. *EPBC Act list of threatened fauna*. Australian Government, Department of the Environment, Canberra, ACT. Available at: http://www.environment.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl#other_animals_vulnerable (accessed 13 May 2014).
- Department of the Environment. 2014b. *Maps: Australia's bioregions (IBRA)*. Department of the Environment, Canberra, ACT. Available at: <http://www.environment.gov.au/topics/land/national-reserve-system/science-maps-and-data/australias-bioregions-ibra> (accessed 24 April 2014).
- Department of the Environment. 2015. *Protected matters search tool*. Australian Government Department of Sustainability, Environment, Water, Population and Communities, Canberra, ACT. Available at: <http://www.environment.gov.au/epbc/pmst/index.html>
- Desmond, A., Kendrick, P. & Chant, A. 2001. Gascoyne 3 (GAS3—Augustus subregion). In: May, J. E. & McKenzie, N. L. (eds) *A biodiversity audit of Western Australia's 53 biogeographical subregions in 2002*. Department of Conservation and Land Management, pp. 240–251.
- DEWHA. 2008a. *Rangelands 2008 - taking the pulse. Gascoyne bioregion*. Department of Environment, Water, Heritage and the Arts, Parkes, ACT. Available at: <http://www.environment.gov.au/system/files/resources/a8015c25-4aa2-4833-ad9c-e98d09e2ab52/files/bioregion-gascoyne.pdf> (accessed 16 April 2015).
- DEWHA. 2008b. *Rangelands 2008 - taking the pulse. Little Sandy Desert bioregion*. Department of Environment, Water, Heritage and the Arts, Parkes, ACT. Available at: <http://www.environment.gov.au/land/publications/acris/pubs/bioregion-little-sandy-desert.pdf> (accessed 25 July 2013).
- DPaW. 2015a. *NatureMap*. Department of Parks and Wildlife, Perth, WA. Available at: <http://naturemap.dec.wa.gov.au/>
- DPaW. 2015b. *Priority Ecological Communities for Western Australia version 21*. Department of Parks and Wildlife, Perth, WA. Available at: http://www.dpaw.wa.gov.au/images/documents/plants-animals/threatened-species/tecs/Priority_ecological_community_list_20_May2014.pdf (accessed 26 November 2015).
- DPaW. 2015c. *Threatened Flora, Fauna and Ecological Communities database searches*. Department of Parks and Wildlife, Kensington, WA. Available at: http://www.dpaw.wa.gov.au/images/documents/plants-animals/threatened-species/Database_Search_request_information_sheet_2015.pdf
- DPaW. 2016a. *List of Threatened Ecological Communities (TECs) endorsed by the Western Australian Minister for the Environment (correct to 6 October 2016)*. Department of Parks and Wildlife, Species and Communities Branch, Perth, WA. Available at: https://www.dpaw.wa.gov.au/images/plants-animals/threatened-species/threatened_ecological_communities_endorsed_by_the_minister_october_2016.pdf (accessed 27 April 2017).
- DPaW. 2016b. *Priority Ecological Communities for Western Australia version 26*. Department of Parks and Wildlife, Perth, WA. Available at: <https://www.dpaw.wa.gov.au/images/plants->

- [animals/threatened-species/priority_ecological_community_list_wa_version_26.pdf](#) (accessed 27 April 2017).
- DPaW. 2017. *Threatened and Priority fauna list*. Department of Parks and Wildlife, Perth, WA. Available at: <https://www.dpaw.wa.gov.au/plants-and-animals/threatened-species-and-communities/threatened-animals> (accessed 6 February 2017).
- Eberhard, S. M., Halse, S. A., Williams, M. R., Scanlon, M. D., Cocking, J. & Barron, H. J. 2009. Exploring the relationship between sampling efficiency and short-range endemism for groundwater fauna in the Pilbara region, Western Australia. *Freshwater Biology* **54**: 885–901.
- Edward, K. L. & Harvey, M. S. 2008. Short-range endemism in hypogean environments: the pseudoscorpion genera *Tyrannochthonius* and *Lagynochthonius* (Pseudoscorpiones: Chthoniidae) in the semiarid zone of Western Australia. *Invertebrate Systematics* **22**: 259–293.
- EPA. 2016a. *Environmental Factor Guideline: Subterranean fauna*. Environmental Protection Authority, Perth, WA. Available at: http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Guideline-Subterranean-Fauna-131216_3.pdf (accessed 20 December 2016).
- EPA. 2016b. *Statement of environmental principles, factors and objectives*. Environmental Protection Authority, Perth, WA. Available at: http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Statement%20of%20Environmental%20Principles%2C%20factors%20and%20objectives_Dec16_1.pdf (accessed 20 December 2016).
- EPA. 2016c. *Technical Guidance: Sampling methods for subterranean fauna*. Environmental Protection Authority, Perth, WA. Available at: http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Tech%20guidance-%20Sampling-Subt-fauna-Dec-2016.pdf (accessed 20 December 2016).
- EPA. 2016d. *Technical Guidance: Sampling of short range endemic invertebrate fauna*. Environmental Protection Authority, Perth, WA. Available at: http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Tech%20guidance-%20Sampling-SREs-Dec-2016.pdf (accessed 20 December 2016).
- EPA. 2016e. *Technical Guidance: Terrestrial fauna surveys*. Environmental Protection Authority, Perth, WA. Available at: http://www.epa.wa.gov.au/sites/default/files/Policies_and_Guidance/Tech%20guidance-%20Terrestrial%20Fauna%20Surveys-Dec-2016.pdf (accessed 10 December 2016).
- Gentili, J. 1979. *Western Landscapes*. University of Western Australia Press, Perth, WA.
- Gray, M. R. 1992. The troglobitic spider genus *Tartarus* Gray with a cladistic analysis of *Tartarus* and *Baiami* Lehtinen (Araneae: Stiphidiidae). *Proceedings of the Linnean Society of New South Wales* **113**: 165–173.
- Grey, K., Hocking, R. M., Stevens, M. K., Bagas, L., Carlsen, G. M., Irimies, F., Pirajno, F., Haines, P. W. & Apak, S. N. 2005. Lithostratigraphic nomenclature of the Officer Basin and correlative parts of the Paterson Orogen Western Australia. *Geological Survey of Australia Report* **93**: 1–89.
- Guzik, M. T., Abrams, K. M., Cooper, S. J. B., Humphreys, W. F., Cho, J. L. & Austin, A. D. 2008. Phylogeography of the ancient Parabathynellidae (Crustacea: Bathynellacea) from the Yilgarn region of Western Australia. *Invertebrate Systematics* **22**: 205–216.
- Hamilton-Smith, E. & Eberhard, S. M. 2000. Conservation of cave communities in Australia. In: Wilkens, H., Culver, D. C. & Humphreys, W. F. (eds) *Ecosystems of the World Vol. 30 - Subterranean ecosystems*. Elsevier, Amsterdam, pp. 647–664.
- Hammer, U. T. 1986. *Saline lake ecosystems of the world*. Dr. W. Junk Publications, Dordrecht, Netherlands.
- Harvey, M. S. 2001a. New cave-dwelling schizomids (Schizomida: Hubbardiidae) from Australia. *Records of the Western Australian Museum, Supplement* **64**: 171–185.

- hr/>
- Harvey, M. S. 2001b. Notes on the spider genus *Symphytognatha* (Araneae: Symphytognathidae) in Western Australia. *Records of the Western Australian Museum* **20**: 345–347.
- Harvey, M. S. 2002. Short-range endemism among the Australian fauna: some examples from non-marine environments. *Invertebrate Systematics* **16**: 555–570.
- Harvey, M. S., Berry, O., Edward, K. L. & Humphreys, G. 2008. Molecular and morphological systematics of hypogean schizomids (Schizomida: Hubbardiidae) in semiarid Australia. *Invertebrate Systematics* **22**: 167–194.
- Harvey, M. S. & Edward, K. L. 2007. A review of the pseudoscorpion genus *Ideoblothrus* (Pseudoscorpiones, Syarinidae) from western and northern Australia. *Journal of Natural History* **41**: 445–472.
- Harvey, M. S., Gray, M. R., Hunt, G. S. & Lee, D. C. 1993. The cavernicolous Arachnida and Myriapoda of Cape Range, Western Australia. In: Humphreys, W. F. (ed.) *The Biogeography of Cape Range, Western Australia*. Western Australian Museum, Perth, WA, pp. 129–144.
- Harvey, M. S. & Humphreys, W. F. 1995. Notes on the genus *Draculoides* Harvey (Schizomida: Hubbardiidae), with the description of a new troglobitic species. *Records of the Western Australian Museum, Supplement* **52**: 183–189.
- Harvey, M. S. & Leng, M. C. 2008. Further observations on *Ideoblothrus* (Pseudoscorpiones: Syarinidae) from subterranean environments in Australia. *Records of the Western Australian Museum* **24**: 381–386.
- Harvey, M. S. & Volschenk, E. S. 2007. Systematics of the Gondwanan pseudoscorpion family Hyidae (Pseudoscorpiones: Neobisioidea); new data and a revised phylogenetic hypothesis. *Invertebrate Systematics* **21**: 365–406.
- Hebert, P. D. N., Ratnasingham, S. & de Waard, J. R. 2003. Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society London B, Supplement* **270**: 96–99.
- Hoch, H. 1993. A new troglobitic planthopper species (Hemiptera: Fulgoroidea: Meenoplidae) from Western Australia. *Records of the Western Australian Museum* **16**: 393–398.
- Hoffman, R. L. 1994. Studies on spirobolid millipedes. XVIII. *Speleostrophus nesiotus*, the first known troglobitic spirobolid millipede, from Barrow Island, Western Australia (Diplopoda, Pachybolidae, Trigoniulinae). *Myriapodologica* **3**: 19–24.
- Holsinger, J. R. 1993. Biodiversity of subterranean amphipod crustaceans: global patterns and zoogeographic implications. *Journal of Natural History* **27**: 821–835.
- Holsinger, J. R. 2000. Ecological derivation, colonization and speciation. In: Wilkens, H., Culver, D. C. & Humphreys, W. F. (eds) *Ecosystems of the World Vol. 30 - Subterranean ecosystems*. Elsevier, Amsterdam, pp. 399–415.
- Hong, S. J. & Cho, J.-L. 2009. Three new species of *Billibathynella* from Western Australia (Crustacea, Syncarida, Parabathynellidae). *Journal of Natural History* **43**: 2365–2390.
- Howarth, F. G. 1983. Ecology of cave arthropods. *Annual Review of Entomology* **28**: 365–389.
- Howarth, F. G. 1993. High-stress subterranean habitats and evolutionary change in cave-inhabiting arthropods. *American Naturalist, Supplement* **142**: 565–577.
- Humphreys, W. F. 1991. Experimental re-establishment of pulse-driven populations in a terrestrial troglobite community. *Journal of Animal Ecology* **60**: 609–623.
- Humphreys, W. F. 1999. Physico-chemical profile and energy fixation in Bundera Sinkhole, an anchialine remiped habitat in north-western Australia. *Journal of the Royal Society of Western Australia* **82**: 89–98.
- Humphreys, W. F. 2000. Background and glossary. In: Wilkens, H., Culver, D. C. & Humphreys, W. F. (eds) *Ecosystems of the World Vol. 30 - Subterranean ecosystems*. Elsevier, Amsterdam, pp. 3–14.

- Humphreys, W. F. 2001. Groundwater calcrete aquifers in the Australian arid zone: the context to an unfolding plethora of stygal biodiversity. *Records of the Western Australian Museum Supplement No. 64*: 63–83.
- Humphreys, W. F. 2008. Rising from Down Under: developments in subterranean biodiversity in Australia from a groundwater fauna perspective. *Invertebrate Systematics* **22**: 85–101.
- Humphreys, W. F. & Shear, W. A. 1993. Troglobitic millipedes (Diplopoda: Paradoxosomatidae) from semi-arid Cape Range, Western Australia: systematics and biology. *Invertebrate Taxonomy* **7**: 173–195.
- Humphreys, W. F., Watts, C. H. S., Cooper, S. J. B. & Leijes, R. 2009. Groundwater estuaries of salt lakes: buried pools of endemic biodiversity on the western plateau, Australia. *Hydrobiologia* **626**: 79–95.
- Jaume, D., Boxshall, G. A. & Humphreys, W. F. 2001. New stygobiont copepods (Calanoida; Misophrioida) from Bundera sinkhole, an anchialine cenote on north-western Australia. *Zoological Journal of the Linnean Society* **133**: 1–24.
- Jaume, D. & Humphreys, W. F. 2001. A new genus of epacteriscid calanoid copepod from an anchialine sinkhole on northwestern Australia. *Journal of Crustacean Biology* **21**: 157–169.
- Javidkar, M., Cooper, S. J. B., King, R. A., Humphreys, W. F. & Austin, A. D. 2015. Molecular phylogenetic analyses reveal a new southern hemisphere oniscidean family (Crustacea : Isopoda) with a unique water transport system. *Invertebrate Systematics* **29**: 554–577.
- Javidkar, M., Cooper, S. J. B., King, R. A., Humphreys, W. F., Bertozzi, T., Stevens, M. I. & Austin, A. D. 2016. Molecular systematics and biodiversity of oniscidean isopods in the groundwater calcretes of Central Western Australia. *Molecular Phylogenetics and Evolution* **104**: 83–98.
- Javidkar, M., King, R. A., Cooper, S. J. B., Humphreys, W. F. & Austin, A. D. 2017. Taxonomy of *Paraplatyarthus* Javidkar and King (Isopoda: Oniscidea: Paraplatyarthridae) with description of five new species from Western Australia, and comments on Australian *Trichorhina* Budde-Lunde, 1908 (Platyarthridae). *Zootaxa* **4243**: 401–431.
- Johnson, S. L., Commander, D. P. & O'Boy, C. A. 1999. *Groundwater resources of the northern Goldfields, Western Australia*. Water and Rivers Commission, East Perth, WA.
- Karanovic, T. 2006. Subterranean copepods (Crustacea, Copepoda) from the Pilbara Region in Western Australia. *Records of the Western Australian Museum, Supplement* **70**: 1–239.
- Knott, B. 1993. Stygofauna from Cape Range Peninsula, Western Australia: Tethyan relicts. *Records of the Western Australian Museum, Supplement* **45**: 109–127.
- Main, B. Y. & Gray, M. R. 1985. Further studies on the systematics of Australian Diplurinae (Chelicerata: Mygalomorphae: Dipluridae): description of the male of *Troglodiplura lowryi*, with notes on its affinities. *Psyche* **92**: 151–162.
- Maly, E. J., Halse, S. A. & Maly, M. P. 1997. Distribution and incidence patterns of *Boeckella*, *Calamoecia*, and *Hemiboeckella* (Copepoda : Calanoida) in Western Australia. *Marine and Freshwater Research* **48**: 615–621.
- Morgan, K. H. 1993. Development, sedimentation and economic potential of palaeoriver systems of the Yilgarn Craton of Western Australia. *Sedimentary Geology* **85**: 637–656.
- Pesce, G. L., De Laurentiis, P. & Humphreys, W. F. 1996. Copepods from ground waters of Western Australia, II. The genus *Halicyclops* (Crustacea: Copepoda: Cyclopidae). *Records of the Western Australian Museum* **18**: 77–85.
- Phoenix. 2014. *Subterranean fauna desktop review of the Lake Disappointment Potash Project*. Phoenix Environmental Sciences Pty Ltd, Balcatta, WA. Unpublished report prepared for Botanica Consulting Pty Ltd on behalf of Reward Minerals Ltd.
- Platnick, N. I. 2008. A new subterranean ground spider genus from Western Australia (Araneae: Trochanteriidae). *Invertebrate Systematics* **22**: 295–299.

-
- Ponder, W. F. & Colgan, D. J. 2002. What makes a narrow-range taxon? Insights from Australian freshwater snails. *Invertebrate Systematics* **16**: 571–582.
- Poulson, T. L. & Lavoie, K. H. 2000. The trophic basis of subsurface ecosystems. In: Wilkens, H., Culver, D. C. & Humphreys, W. F. (eds) *Ecosystems of the World Vol. 30 - Subterranean ecosystems*. Elsevier, Amsterdam, pp. 231–250.
- Preston Consulting. 2017. *Beyondie Sulphate of Potash Project. Section 38 referral - supporting information*. Preston Consulting Pty Ltd, East Perth, WA.
- Roth, L. M. 1991. A new cave-dwelling cockroach from Western Australia (Blattaria: Nocticolidae). *Records of the Western Australian Museum* **15**: 17–21.
- Shear, W. A. & Humphreys, W. F. 1996. A new *Stygiochiropus* from a North West Cape (Western Australia) coastal plain cave (Diplopoda: Polydesmida: Paradoxosomatidae). *Records of the Western Australian Museum* **17**: 447–449.
- Stoch, F. 2001. How many species of *Diacyclops*? New taxonomic characters and species richness in a freshwater cyclopoid genus (Copepoda, Cyclopoida). *Hydrobiologia* **453/454**: 525–531.
- Subterranean Ecology. 2010a. *Gold Fields, St Ives Gold Mines troglotauna desktop assessment*. Subterranean Ecology (Scientific Environmental Services) Pty Ltd, Stirling, WA. Unpublished report prepared for St Ives Gold Mines.
- Subterranean Ecology. 2010b. *Gold Fields, St Ives Gold Mines. Stygofauna desktop assessment*. Subterranean Ecology Pty Ltd, Stirling, WA. Unpublished report prepared for Gold Fields Pty Ltd.
- Thackway, R. & Cresswell, I. D. 1995. *An Interim Biogeographic Regionalisation for Australia: a framework for establishing the national system of reserves, Version 4.0*. Australian Nature Conservation Agency, Canberra, ACT.
- Toro Energy. 2011. *Wiluna Uranium Project. Appendix F: Technical reports - Salt lake ecology and subterranean fauna*. Toro Energy Ltd, West Perth, WA. Public Environmental Review.
- Toro Energy. 2012. *Appendix E - Subterranean fauna (stygofauna)*. Toro Energy Ltd, Perth, WA. Memo - Wiluna Uranium Project Stygofauna Fauna April, 2012.
- van Leeuwen, S. 2002. *Biological survey of the south-western Little Sandy Desert*. Department of Conservation and Land Management, Perth, WA. Final report, National Reserve System Project N706.
- Vandel, A. 1974 [imprint date 1973]. Les isopodes terrestres de l'Australie. Étude systématique et biogéographique. *Mémoires de Muséum national d'Histoire naturelle, Paris* **82**: 1–171.
- Volschenk, E. S. & Prendini, L. 2008. *Aops oncodactylus*, gen. et sp. nov., the first troglobitic urodacid (Urodacidae: Scorpiones), with a re-assessment of cavernicolous, troglobitic and troglomorphic scorpions. *Invertebrate Systematics* **22**: 235–257.
- Watts, C. H. S. & Humphreys, W. F. 2001. A new genus and six species of Dytiscidae (Coleoptera) from underground waters in the Yilgarn paleodrainage system of Western Australia. *Records of the South Australian Museum* **34**: 99–114.
- Watts, C. H. S. & Humphreys, W. F. 2004. Thirteen new Dytiscidae (Coleoptera) of the genera *Boongurrus* Larson, *Tjirtudessus* Watts & Humphreys and *Nirripiriti* Watts and Humphreys, from underground waters in Australia. *Transactions of the Royal Society of South Australia* **128**: 99–129.
- Western Australian Government. 2015. Wildlife Conservation Act 1950 - Wildlife Conservation (Specially Protected Fauna) Notice 2015. *Western Australian Government Gazette* **166**: 4532–4543.
- Wilson, R. S. & Humphreys, W. F. 2001. *Prionospio thalanji* sp. nov. (Polychaeta: Spionidae) from an anchialine cave, Cape Range, north-west Western Australia. *Records of the Western Australian Museum Supplement No. 64*: 105–113.

Yager, J. & Humphreys, W. F. 1996. *Lasionectes exleyi*, sp. nov., the first remipede crustacean recorded from Australia and the Indian Ocean, with a key to the world species. *Invertebrate Taxonomy* **10**: 171–187.

Appendix 1 Conservation significant subterranean fauna in Western Australia (DPaW 2017)

Genus and species	Common name	Trogl-/ stygo fauna	Conservation rating (DPaW 2017)	Distribution (DPaW 2017)	Currently known records
Arachnida					
Araneae (spiders)					
<i>Tartarus mullamullangensis</i>	Mullamullalang Cave spider	T	VU	South Coast	Mullamullang Cave, Nullarbor Plain (Gray 1992)
<i>Tartarus murdochensis</i>	Murdoch Sink cave spider	T	VU	South Coast	Phyllistine Flattener Cave and Murdoch Sink, Nullarbor Plain (Gray 1992)
<i>Tartarus nurinensis</i>	Nurina Cave spider	T	VU	South Coast	Nurina Cave, Roe Plains (Gray 1992)
<i>Tartarus thampannensis</i>	Thampanna Cave spider	T	VU	South Coast	Thampanna Cave, Nullarbor Plain (Gray 1992)
<i>Troglodiplura lowryi</i>	Nullarbor cave trapdoor spider	T	VU	South Coast	Roaches Rest Cave and Cave NR. 6 Bore, Nullarbor Plain (Main & Gray 1985)
Pseudoscorpiones					
<i>Ideoblothrus linnaei</i>		T	P1	Pilbara	Mesa A (Harvey & Leng 2008)
<i>Ideoblothrus</i> sp. Mesa A (WAM T81374)		T	P1	Pilbara	Mesa A (Harvey & Edward 2007)
<i>Indohya damocles</i>	Cameron's Cave pseudoscorpion	T	CR	Pilbara	Cameron's Cave, Cape Range (Harvey & Volschenk 2007)
<i>Lagynochthonius asema</i>		T	P1	Pilbara	Mesa A (Edward & Harvey 2008)
<i>Tyrannochthonius</i> sp. Mesa A (WAM T81480)					Mesa A
Schizomida					
<i>Bamazomus subsolans</i>	Eastern Cape Range bamazomus	T	EN	Pilbara	Unnamed limestone quarry, Cape Range (Harvey 2001a)
<i>Bamazomus vespertinus</i>	western Cape Range bamazomus	T	EN	Pilbara	Cave C-215, Cape Range Peninsula (Harvey 2001a)

Genus and species	Common name	Trogl-/ stygo fauna	Conservation rating (DPaW 2017)	Distribution (DPaW 2017)	Currently known records
<i>Draculoides bramstokeri</i>	Barrow Island draculoides	T	VU	Pilbara	Barrow Island (Harvey & Humphreys 1995)
<i>Draculoides brooksi</i>	northern Cape Range draculoides	T	EN	Pilbara	North-eastern Cape Range Peninsula (Harvey <i>et al.</i> 2008)
<i>Draculoides julianneae</i>	western Cape Range draculoides	T	EN	Pilbara	Caves C-111 and C-215, Cape Range Peninsula (Harvey <i>et al.</i> 2008)
<i>Draculoides mesozeirus</i>	Middle Robe draculoides	T	VU	Pilbara	Middle Robe (Harvey <i>et al.</i> 2008)
<i>Draculoides vinei</i>	Cape Range Draculoides	T	P4	Pilbara	Caves within Tulki limestone of Cape Range (Harvey 2001a; Harvey <i>et al.</i> 2008)
<i>Paradraculoides anachoretus</i>	Mesa A paradraculoides	T	VU	Pilbara	Mesa A (Harvey <i>et al.</i> 2008)
<i>Paradraculoides bythius</i>	Mesa B/C paradraculoides	T	VU	Pilbara	Mesa B and Mesa C (Harvey <i>et al.</i> 2008)
<i>Paradraculoides gnophicola</i>	Mesa G paradraculoides	T	VU	Pilbara	Mesa G (Harvey <i>et al.</i> 2008)
<i>Paradraculoides kryptus</i>	Mesa K paradraculoides	T	VU	Pilbara	Mesa K (Harvey <i>et al.</i> 2008)
Myriapoda					
<i>Speleostrophus nesiotes</i>	Barrow Island millipede	T	VU	Pilbara	Barrow Island (Car <i>et al.</i> 2013; Hoffman 1994)
<i>Stygiochiropus peculiaris</i>	Cameron's Cave millipede	T	CR	Pilbara	Camerons Cave, Cape Range Peninsula (Shear & Humphreys 1996)
<i>Stygiochiropus isolatus</i>	Millipede	T	VU	Pilbara	Cave C-222, Cape Range Peninsula (Humphreys & Shear 1993)
<i>Stygiochiropus sympatricus</i>	Millipede	T	VU	Pilbara	Cave C-111, Cape Range Peninsula (Humphreys & Shear 1993)
Crustacea					
Amphipoda					
<i>Bogidomma australis</i>	Barrow Island bogidomma amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)

Genus and species	Common name	Troglo-/ stygo fauna	Conservation rating (DPaW 2017)	Distribution (DPaW 2017)	Currently known records
<i>Hurleya</i> sp. (WAM 642–97)	Crystal Cave crangonyctoid amphipod	S	CR	Swan	Crystal Cave, Yanchep
<i>Liagoceradocus branchialis</i>	Cape Range liagoceradocus amphipod	S	EN	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Bradbury & Williams 1996b)
<i>Liagoceradocus subthalassicus</i>	Barrow Island liagoceradocus amphipod	S	VU	Pilbara	Ledge Cave B-1, Barrow Island (Bradbury & Williams 1996b)
<i>Nedsia fragilis</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia humphreysi</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia hurlberti</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia macrosculptilis</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia sculptilis</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia straskraba</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia urifimbriata</i>	Freshwater amphipod	S	VU	Pilbara	Barrow Island (Bradbury & Williams 1996a)
<i>Nedsia chevronia</i>	Freshwater amphipod	S	P2	Pilbara	Barrow Island (Bradbury 2002)
Copepoda					
<i>Bunderia misophaga</i>	Copepod	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Jaume & Humphreys 2001)
<i>Speleophria bunderae</i>	Copepod	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Jaume <i>et al.</i> 2001)
<i>Stygocyclopia australis</i>	Copepod	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Jaume <i>et al.</i> 2001)
Decapoda					
<i>Stygiocaris lancifera</i>	Lance-beaked cave shrimp	S	VU	Pilbara	Cape Range Peninsula (Knott 1993)
<i>Stygiocaris stylifera</i>	Spear-beaked Cave Shrimp	S	P4	Pilbara	Cape Range Peninsula (Knott 1993)

Genus and species	Common name	Trogl-/ stygo fauna	Conservation rating (DPaW 2017)	Distribution (DPaW 2017)	Currently known records
Isopoda					
<i>Abebaioscia troglodytes</i>	Pannikin Plain Cave isopod	T	VU	South Coast	Pannikin Plain Cave, Nullarbor Plain (Vandel 1974 [imprint date 1973])
<i>Paraplatyarthus subterraneus</i>	Poseidon slater	T	P1		Pilbara (Javidkar & King 2015)
Ostracoda					
<i>Welesina kornickeri</i>	Ostracod	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Danielopol <i>et al.</i> 2000)
Remipedia					
<i>Kumonga exleyi</i>	Cape Range remiped	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Yager & Humphreys 1996)
Polychaeta					
<i>Prionospio thalANJI</i>	Bristle worm	S	CR	Pilbara	Bundera Sinkhole, Cape Range Peninsula (Wilson & Humphreys 2001)
Insecta					
Blattaria					
<i>Nocticola flabella</i>	Cape Range Blind Cockroach	T	P2	Pilbara	Cape Range Peninsula (Roth 1991)

Appendix 2 Subterranean Threatened Ecological Communities in Western Australia (DPaW 2016a)

Name of community	Description	Category of Threat and criteria met under WA criteria	Conservation Rating (EPBC Act 1999)	
Caves SP01	Aquatic Root Mat Community Number 1 of Caves of the Swan Coastal Plain	CR B) i), CR B) ii)	EN	Swan Coastal Plain
Caves Leeuwin01	Aquatic Root Mat Community Number 1 of Caves of the Leeuwin Naturaliste Ridge	CR B) i), CR B) ii)	EN	Warren
Caves Leeuwin02	Aquatic Root Mat Community Number 2 of Caves of the Leeuwin Naturaliste Ridge	CR B) i), CR B) ii)	EN	Warren
Caves Leeuwin03	Aquatic Root Mat Community Number 3 of Caves of the Leeuwin Naturaliste Ridge	CR B) i), CR B) ii)	EN	Warren
Caves Leeuwin04	Aquatic Root Mat Community Number 4 of Caves of the Leeuwin Naturaliste Ridge	CR B) i), CR B) ii)	EN	Warren
Cameron's	Cameron's Cave Troglobitic Community	CR B) i), CR B) ii)		Carnarvon Basin
Bundera	Cape Range Remiped Community	CR B) ii)		Carnarvon Basin
Ethel Gorge	Ethel Gorge aquifer stygobiont community	EN B) ii)		Pilbara
Depot Springs	Depot Springs stygofauna community	VU B)		Goldfields Region, Murchison Bioregion

Appendix 3 Priority Ecological Communities (DPaW 2016b)

Name of community	Description	Threats	Category (WA)
Pilbara			
Barrow Island subterranean fauna	Barrow Island stygofauna and troglotauna	Mining and industrial development	Priority 1
Subterranean invertebrate communities of mesas in the Robe Valley region	A series of isolated mesas occur in the Robe Valley in the state's Pilbara region. The mesas are remnants of old valley infill deposits of the palaeo Robe River. The troglotic faunal communities occur in an extremely specialised habitat and appear to require the particular structure and hydrogeology associated with mesas to provide a suitable humid habitat. Short-range endemism is common in the fauna. The habitat is the humidified pisolitic strata	Mining	Priority 1
Subterranean invertebrate community of pisolitic hills in the Pilbara	A series of isolated low undulating hills occur in the state's Pilbara region. The troglotauna are being identified as having very short-range distributions	Mining	Priority 1
Mingah Springs calcrete groundwater assemblage type on Gascoyne palaeodrainage on Mingah Spring Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Stygofaunal community of the Bungaroo Aquifer	A unique assemblage of aquatic subterranean fauna including eels, snails and other stygofauna	Groundwater drawdown, mining	Priority 1
Stygofaunal communities of the Western Fortescue Plains freshwater aquifer	A unique assemblage of subterranean invertebrate fauna	Groundwater drawdown and salinisation	Priority 4(ii)
Kimberley			
Invertebrate community of Napier Range Cave	On Old Napier Downs, Karst No. KNI	Mine close by and tourist visitation	Priority 1
Midwest			
Badja calcrete groundwater assemblage type on Moore palaeodrainage on Badja Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Belele calcrete groundwater assemblage type on Murchison palaeodrainage on Belele Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Beringarra calcrete groundwater assemblage type on Murchison palaeodrainage on Beringarra Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Black Range South and Windsor groundwater calcrete assemblage type on Raeside and Murchison palaeodrainage on Lake Mason and Windsor Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Bunnawarra calcrete groundwater assemblage type on Moore palaeodrainage on Bunnawarra Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Byro Central and Byro HS calcrete groundwater assemblage types on Murchison palaeodrainage on Byro Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Challa, Challa North and Wondinong calcrete groundwater assemblage type on Murchison palaeodrainage on Challa and Wondinong Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Cogla Downs calcrete groundwater assemblage type on Murchison palaeodrainage on Yarrabubba Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Curbur calcrete groundwater assemblage type on Gascoyne palaeodrainage on Curbur Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Dalgety and Landor calcrete groundwater assemblage type on Gascoyne palaeodrainage on Dalgety Downs and Landor Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Doolgunna calcrete groundwater assemblage type on Gascoyne palaeodrainage on Doolgunna Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Gabyon calcrete groundwater assemblage type on Moore palaeodrainage on Gabyon Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Gifford Creek, Mangaroon, Wanna calcrete groundwater assemblage type on Lyons palaeodrainage on Gifford Creek, Lyons and Wanna Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Hillview calcrete groundwater assemblage type on Murchison palaeodrainage on Hillview Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Innouendy calcrete groundwater assemblage type on Murchison palaeodrainage on Innouendy Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Karalundi calcrete groundwater assemblage type on Murchison palaeodrainage on Karalundi Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Killara calcrete groundwater assemblage types on Murchison palaeodrainage on Killara Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Killara North calcrete groundwater assemblage types on Murchison palaeodrainage on Killara Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Austin calcrete groundwater assemblage type on Murchison palaeodrainage on Austin Downs Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Maranalgo west calcrete assemblage type on Moore palaeodrainage on Maranalgo Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Meeberrie calcrete groundwater assemblage type on Murchison palaeodrainage on Meeberrie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Meka calcrete groundwater assemblage type on Murchison palaeodrainage on Meka Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Milgun central calcrete groundwater assemblage types on Gascoyne palaeodrainage on Milgun Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Milgun south calcrete groundwater assemblage types on Gascoyne palaeodrainage on Milgun Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Milly Milly calcrete groundwater assemblage type on Murchison palaeodrainage on Milly Milly Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Mount Augustus calcrete groundwater assemblage type on Lyons palaeodrainage on Mount Augustus Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Mt Clere calcrete groundwater assemblage type on Gascoyne palaeodrainage on Mt Clere Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Mount Narryer calcrete groundwater assemblage type on Murchison palaeodrainage on Mount Narryer Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Mount Padbury calcrete groundwater assemblage type on Murchison palaeodrainage on Mount Padbury Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Muralgarra calcrete groundwater assemblage type on Murchison palaeodrainage on Muralgarra Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Murchison Downs calcrete groundwater assemblage type on Murchison palaeodrainage on Murchison Downs Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Ninghan calcrete groundwater assemblage type on Moore palaeodrainage on Ninghan Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Nowthanna Hill calcrete groundwater assemblage type on Murchison palaeodrainage on Yarrabubba Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Paroo calcrete groundwater assemblage type on Carey palaeodrainage on Paroo Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Polelle calcrete groundwater assemblage type on Murchison palaeodrainage on Polelle Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Taincrow calcrete groundwater assemblage type on Murchison palaeodrainage on Taincrow Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Three Rivers calcrete groundwater assemblage types on Gascoyne palaeodrainage on Three Rivers Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Three Rivers Plutonic calcrete groundwater assemblage types on Gascoyne palaeodrainage on Three Rivers Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Wagga Wagga and Yalgoo calcrete groundwater assemblage type on Yalgoo and Moore palaeodrainage on Wagga Wagga and Bunnawarra Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Windimurra calcrete groundwater assemblage type on Murchison palaeodrainage on Windimurra Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Wooramel calcrete groundwater assemblage type on Wooramel palaeodrainage on Innouendy Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yarrabubba east calcrete groundwater assemblage types on Murchison palaeodrainage on Yarrabubba Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yarrabubba west calcrete groundwater assemblage types on Murchison palaeodrainage on Yarrabubba Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Yoweragabbie calcrete groundwater assemblage type on Moore palaeodrainage on Yoweragabbie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Goldfields			
Albion Downs calcrete groundwater assemblage type on Carey palaeodrainage on Albion Downs Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Banjawarn and Melrose (Lake Darlot) calcrete groundwater assemblage type on Carey palaeodrainage on Banjawarn and Melrose Stations	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Barwidgee calcrete groundwater assemblage type on Carey palaeodrainage on Barwidgee Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Black Range North calcrete groundwater assemblage type on Raeside palaeodrainage on Lake Mason Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Cunyu SBF and Cunyu Sweetwater calcrete groundwater assemblage types on Nabberu palaeodrainage on Cunyu Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Dandaraga calcrete groundwater assemblage type on Raeside palaeodrainage on Dandaraga Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Glenayle and Carnegie Downs calcrete groundwater assemblage type on Burnside palaeodrainage on Glenayle Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Hinkler Well calcrete groundwater assemblage type on Carey palaeodrainage on Lake Way Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Way South calcrete groundwater assemblage type on Carey palaeodrainage on Lake Way Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Jundee Homestead calcrete groundwater assemblage type on Carnegie palaeodrainage on Jundee Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Jundee South Hill calcrete groundwater assemblage type on Carnegie palaeodrainage on Jundee Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Kaluwiri calcrete groundwater assemblage type on Raeside palaeodrainage on Kaluwiri Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Mason calcrete groundwater assemblage type on Raeside palaeodrainage on Lake Mason Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Miranda east calcrete groundwater assemblage types on Carey palaeodrainage on Yakabindie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Miranda west calcrete groundwater assemblage types on Carey palaeodrainage on Yakabindie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lake Violet south and Lake Violet calcrete groundwater assemblage types on Carey palaeodrainage on Millbillillie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Laverton Downs calcrete groundwater assemblage type on Carey palaeodrainage on Laverton Downs Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Lorna Glen calcrete groundwater assemblage type on Carnegie palaeodrainage on Lorna Glen Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Melita calcrete groundwater assemblage type on Raeside palaeodrainage on Melita Station (Sons of Gwalia)	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Millbillillie: Bubble calcrete groundwater assemblage type on Carey palaeodrainage on Millbillillie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Mount Morgan calcrete groundwater assemblage type on Carey palaeodrainage on Mount Weld Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Nambi calcrete groundwater assemblage type on Carey palaeodrainage on Nambi Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Old Cunya calcrete groundwater assemblage type on Nabberu palaeodrainage on Cunyu Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Perrinvale (Pine Well) calcrete groundwater assemblage type on Raeside palaeodrainage on Perrinvale Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Pinnacles calcrete groundwater assemblage type on Raeside palaeodrainage on Pinnacles Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1

Name of community	Description	Threats	Category (WA)
Sturt Meadows calcrete groundwater assemblage type on Raeside palaeodrainage on Sturt Meadows Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Uramurdah Lake calcrete groundwater assemblage type on Carey palaeodrainage on Millbillillie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Wiluna BF calcrete groundwater assemblage type on Carey palaeodrainage on Millbillillie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Windidda calcrete groundwater assemblage type on Carnegie palaeodrainage on Windidda Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yakabindie calcrete groundwater assemblage type on Carey palaeodrainage on Yakabindie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yandal calcrete groundwater assemblage type on Carey palaeodrainage on Yandal Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yeelirrie calcrete groundwater assemblage type on Carey palaeodrainage on Yeelirrie Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Yuinmery calcrete groundwater assemblage types on Raeside palaeodrainage on Yuinmery Station	Unique assemblages of invertebrates have been identified in the groundwater calcretes	Mining	Priority 1
Warren			
Microbial mantles of Nullarbor caves (especially Weebubbie Cave)	Significant microbial communities in underwater sections of cave	Uncontrolled access	Priority 1
Subterranean faunal ecosystems of Nullarbor caves (known from Nurina Cave, Olwolgine Cave, Burnabbie Cave, N327, N1327)	The caves contain communities of invertebrates, other fauna and sensitive habitats including tree roots. Caves included in this community contain at least four troglobitic taxa.	Uncontrolled access	Priority 3(i)

