

FLORA AND VEGETATION SURVEY OF THE MANGLES BAY AREA CAPE PERON, ROCKINGHAM

Prepared for

STRATEGEN CONSULTING



Job No: 09.232

Report No: 09/055



Australia

FLORA AND VEGETATION SURVEY OF THE MANGLES BAY AREA CAPE PERON, ROCKINGHAM

Prepared for

STRATEGEN CONSULTING

Prepared by

ENV Australia Pty Ltd
Level 1, 503 Murray Street
PERTH WA 6000
Phone: (08) 9214 6100
Fax: (08) 9226 4109
Email: env@env.net.au

Prepared by:	<i>Narelle Whittington</i>
Status:	<i>Final</i>
QA Review:	<i>Peter Jobson</i>
Technical Review:	<i>Teresa Gepp</i>
Content Review:	<i>Teresa Gepp</i>
Date:	<i>1 February 2011</i>

TABLE OF CONTENTS

EXECUTIVE SUMMARY	III
1 INTRODUCTION	1
1.1 LOCATION.....	1
1.2 CLIMATE.....	2
1.3 REGIONAL SOILS, LANDFORMS AND VEGETATION	2
1.4 PROTECTION OF FLORA AND VEGETATION	4
1.5 INTRODUCED SPECIES.....	6
1.6 BUSH FOREVER.....	6
2 METHODS.....	8
2.1 FLORA SURVEY VARIABLES.....	10
2.2 PERMITS	12
3 RESULTS	13
3.2 FIELD SURVEY - FLORA	13
3.3 FIELD SURVEY – VEGETATION	17
3.3.1 Floristic Community Types	17
3.3.2 Priority Ecological Communities and Threatened Ecological Communities.....	21
3.3.3 Vegetation Condition	21
3.3.4 Bush Forever.....	22
4 DISCUSSION	23
5 SUMMARY AND RECOMMENDATIONS.....	28
5.1 RECOMMENDATIONS.....	29
6 REFERENCES	30

FIGURES

FIGURE 1	SITE LOCATION
FIGURE 2	FLORA SURVEY AREA
FIGURE 3	RAINFALL AND TEMPERTURE
FIGURE 4	QUADRAT AND SIGNIFICANT FLORA LOCATIONS
FIGURE 5	FLORISTIC COMMUNITY TYPE MAP
FIGURE 6	BUSHLAND CONDITION MAP
FIGURE 7	AREA AND CONDITION OF TEC WITHIN THE IMPACT AREA

TABLES

TABLE 1	LIMITATIONS AND CONSTRAINTS ASSOCIATED WITH THE [SITE] FLORA AND VEGETATION SURVEY
TABLE 2	SIGNIFICANT FLORA LOCATIONS
TABLE 3	FLORA POTENTIALLY SENSITIVE TO GROUNDWATER CHANGES
TABLE 4	WEED SPECIES IDENTIFIED
TABLE 5	PRIMER ANANYSIS RESULTS
TABLE 6	FLORISTIC COMMUNITY TYPE DETERMINATION

APPENDICES

APPENDIX A	DEFINITIONS OF DECLARED RARE / PRIORITY / THREATENED FLORA AND THREATENED / PRIORITY ECOLOGICAL COMMUNITIES
APPENDIX B	ENVIRONMENTAL WEEDS AND DECLARED PLANT CATEGORIES
APPENDIX C	DEC DATABASE SEARCH RESULTS
APPENDIX D	MATRIX OF SPECIES FOUND AT EACH SITE
APPENDIX E	FLORA SURVEY FIELD DATASHEETS AND SITE PHOTOGRAPHS
APPENDIX F	BUSH FOREVER CONDITION SCALE
APPENDIX G	FLORISTICS DENDOGRAM

STATEMENT OF LIMITATIONS

Scope of Services

This environmental site assessment report ('the report') has been prepared in accordance with the scope of services set out in the contract, or as otherwise agreed, between the Client and ENV.Australia Pty Ltd (ENV) ('scope of services'). In some circumstances the scope of services may have been limited by factors such as time, budget, access and/or site disturbance constraints.

Reliance on Data

In preparing the report, ENV has relied on data, surveys, analyses, designs, plans and other information provided by the Client and other individuals and organisations, most of which are referred to in the report ('the data'). Except as otherwise stated in the report, ENV has not verified the accuracy or completeness of the data. To the extent that the statements, opinions, facts, information, conclusions and/or recommendations in the report ("conclusions") are based in whole or in part on the data, those conclusions are contingent upon the accuracy and completeness of the data. ENV will not be liable in relation to incorrect conclusions should any data, information or condition be incorrect or have been concealed, withheld, unavailable, misrepresented or otherwise not fully disclosed to ENV.

Environmental Conclusions

In accordance with the scope of services, ENV has relied on the data and has conducted environmental field monitoring and/or testing in the preparation of the report. The nature and extent of monitoring and/or testing conducted is described in the report.

Within the limitations imposed by the scope of services, the monitoring, testing, sampling and preparation of this report have been undertaken and performed in a professional manner, in accordance with generally accepted practices and using a degree of skill and care ordinarily exercised by reputable environmental consultants under similar circumstances. No other warranty, express or implied, is made.

Report for Benefit of Client

The report has been prepared for the benefit of the Client and for no other party. ENV assumes no responsibility and will not be liable to any other person or organisation for or in relation to any matter dealt with or conclusions expressed in the report, or for any loss or damage suffered by any other person or organisation arising from matters dealt with or conclusions expressed in the report (including, without limitation, matters arising from any negligent act or omission of ENV or for any loss or damage suffered by any other party relying on the matters dealt with or conclusions expressed in the report). Other parties should not rely upon the report or the accuracy or completeness of any conclusions, and should make their own enquiries and obtain independent advice in relation to such matters.

Other Limitations

ENV will not be liable to update or revise the report to take into account any events or circumstances occurring or facts becoming apparent after the date of the report.

The scope of services did not include any assessment of the title to or ownership of the properties, buildings and structures referred to in the report, nor the application or interpretation of laws in the jurisdiction in which those properties, buildings and structures are located.

EXECUTIVE SUMMARY

ENV.Australia Pty Ltd (ENV) was commissioned by Strategen in October 2009 to undertake a targeted declared rare flora, priority flora and floristic community type assessment for the Mangles Bay Area in Cape Peron, Rockingham (the project area). The assessment has been undertaken as part of the concept planning process for the development of a marina-based tourism precinct.

This targeted spring survey, recommended by Bennett (2005), was undertaken to supplement and complete the botanical data for the survey area. In particular the objectives of the assessment were to: re-assess Floristic Community Types; determine the presence of Threatened Ecological Communities; and re-survey for Declared Rare and Priority Flora species.

A database search of the area resulted in four Declared Rare, 15 Priority Flora species and four Threatened Ecological Communities being identified as potentially occurring in the area.

During the survey, a total of 75 taxa, from 37 families and 65 genera were recorded within the survey area (41 native flora taxa and 34 introduced taxa).

No Endangered species pursuant to the *Environment Protection and Biodiversity Conservation Act 1999*, Declared Rare Flora pursuant to the *Wildlife Conservation Act 1950* or Priority Flora species listed by the Department of Environment and Conservation were located.

One Declared Plant (i.e. weed) species was found in the study area:

- **Asparagus asparagoides* (Bridal Creeper), this species is listed as Priority 1 for the whole State.

The site is mapped as containing the Quindalup Vegetation Complex: Coastal dune complex consisting mainly of two alliances – the standard fore-dune alliance and the mobile and stable dune alliance. Local variations include the low closed forest of *Melaleuca lanceolata* – *Callitris preissii* and the closed scrub of *Acacia rostellifera*. This complex exceeds the 10% retention status recommended for Western Australia by the Environmental Protection Authority's *Position Statement No. 2* and is therefore considered to be adequately represented.

The following Floristic Community Types have been identified as occurring on site:

- **SCP16:** Highly saline seasonal wetlands
- **SCP29a:** Coastal shrublands on shallow sands
- **SCP29b:** *Acacia* shrublands on taller dunes
- **S13:** Northern *Olearia axillaris* – *Scaevola crassifolia* shrublands

- **S14:** *Spinifex longifolius* grasslands and low shrublands
- **SCP30a:** *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands
- **SCP30b:** Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands
- **S15:** Weed Group

One Threatened Ecological Community, Floristic Community Type SCP30a *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands was identified as occurring in three locations within the project area. This Floristic Community Type is listed as Vulnerable by the State but is not listed by the Commonwealth.

The Floristic Community Type 30a is located on the boundary of the school is not a good representation of the Floristic Community Type and is not likely to be a viable site to preserve due to its perimeter to area ratio, surrounding degrading influences and the lack of understorey species.

The Threatened Ecological Community site surrounding the small grassed area on the corner of Memorial Drive and Safety Bay Road has been subject to various degrading influences such as weeds and human activity, however the site is surrounded by native vegetation and therefore has the potential to be remediated to improve its condition. There are many informal tracks that dissect the area and rubbish has been dumped adjacent to the site so these degrading influences will need to be resolved if the vegetation community is to be conserved.

The third Threatened Ecological Community site located at the base of the most western car park has a walking track that runs adjacent to the site and, although it experiences degradation from people deviating off the path, is still in relatively good condition. The spread of weeds needs to be resolved to protect the vegetation community.

Two of the Floristic Community Types are listed as Priority 3 Priority Ecological Communities by the Department of Environment and Conservation; Floristic Community Type SCP30b Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands and Floristic Community Type SCP29b *Acacia* shrublands on taller dunes. Priority Ecological Communities are not protected under legislation however they should be treated as significant and taken into consideration during the planning phase.

1 INTRODUCTION

ENV.Australia Pty Ltd (ENV) was commissioned by Strategen in October 2009 to undertake a targeted Declared Rare Flora, priority flora and floristic community type assessment for the Mangles Bay Area in Cape Peron, Rockingham (the project area). The assessment has been undertaken as part of the concept planning process for the LandCorp Cape Peron marina-based tourist precinct development (the Project).

Bennett Environmental Consulting Pty Ltd previously surveyed the project area in 2005, identifying Floristic Community Types and the potential presence of Threatened Ecological Communities. A flora and vegetation survey was also conducted by Keating and Trudgen in October 1986.

This targeted spring survey, recommended by Bennett (2005), was undertaken to supplement and complete the botanical data for the survey area. In particular the objectives of the assessment were to: re-assess Floristic Community Types; determine the presence of Threatened Ecological Communities; and re-survey for Declared Rare and Priority Flora species.

1.1 LOCATION

The survey area is located approximately 39 kilometres to the south-west of Perth's Central Business District in the Swan Coastal Plain region of Western Australia (Figure 1) and lies within the suburbs of Peron and Shoalwater on the shores of Mangles Bay, Rockingham. The survey area is bounded by Safety Bay Road and Boundary Road (Figure 2) and includes the Core Project Area for the Cape Peron marina-based tourist precinct development (impact area).

The site is located in the southwest province of Western Australia in the Darling Botanical District and within the Swan Coastal Plain Subregion in the Drummond Botanical Subdistrict (Beard 1990). The Drummond Botanical Subdistrict consists mainly of the following vegetation communities:

- *Banksia* Low Woodland on leached sands and *Melaleuca* Swamps in poorly drained areas;
- Woodland of Tuart (*Eucalyptus gomphocephala*); and
- Jarrah (*Eucalyptus marginata*) and Marri (*Corymbia calophylla*) woodlands on the less leached soils (Beard 1990).

1.2 CLIMATE

The climate of this region is warm Mediterranean, with an average maximum summer temperature of 28.3°C and an average minimum winter temperature of 10.9°C (Figure 3; Bureau of Meteorology [BoM] 2009). The region receives an average annual rainfall of 759.3mm, with the majority of precipitation occurring in winter (BoM 2009).

The Perth area received a slightly lower amount of rainfall than average in the three months preceding the survey (August to October), with the area having received 222 mm (Figure 3). On average the area usually receives 235.3 mm of rainfall over this same period. For the two months of spring (September and October) preceding the survey the area received 97.2 mm, compared with 117.4 mm for the long term average for the same period (BoM 2009).




Figure 3: Average Monthly Rainfall and Maximum and Minimum Temperatures at Perth Airport from 1944-2009 (BoM 2009)

1.3 REGIONAL SOILS, LANDFORMS AND VEGETATION

For a development proposal to be assessed in terms of the flora and vegetation values that may be impacted upon, an understanding of the vegetation

communities at the site in question is required. In Western Australia, there are various floristic reports that detail a region's botanical values.

A widely-used vegetation classification system that maps and describes vegetation communities in south-west Western Australia is *Vegetation of the Darling System* in the *Atlas of Natural Resources, Darling System, Western Australia* (Department of Conservation and Environment 1980). This document describes vegetation communities as vegetation complexes, and maps the distribution of each complex.

Vegetation complexes are defined as a combination of distinct site vegetation types, usually associated with a particular geomorphic, climatic, floristic and vegetation structural association. Vegetation complexes are based on the pattern of vegetation at a regional scale, as it reflects the underlying key determining factors of landforms, climate and soils.

The soils and landform unit, as well as the vegetation complex Cape Peron supports, is described below:

1.3.1 Soils and Landforms

The site occurs on the Swan Coastal Plain portion of the Darling System (Churchward and McArthur 1978). The Swan Coastal Plain consists of aeolian and fluvial deposits: specifically the site is on:

- Quindalup Unit: Dunes and beach ridges composed of calcareous sand.

1.3.2 Vegetation Complex Mapping

Hedde *et al.* (1978) mapped the area as containing one Swan Coastal Plain vegetation complex which is related to the underlying soil profile:

- Quindalup Complex: Coastal dune complex consisting mainly of two alliances – the standard fore-dune alliance and the mobile and stable dune alliance. Local variations include the low closed forest of *Melaleuca lanceolata* – *Callitris preissii* and the closed scrub of *Acacia rostellifera*.

The Environmental Protection Authority (EPA) recognises that native vegetation complexes which have less than 10% of pre-European clearing extent remaining in the Bush Forever study area may be considered regionally significant (EPA 2006). Proposals that would impact on a vegetation complex with 10% or less remaining are likely to be formally assessed by the EPA (EPA 2006).

Bush Forever gives an estimate of the percentage of each complex that remains within the Bush Forever study area compared to its pre-European settlement extent, so an estimate of the scarcity of each complex can be determined. On the

Swan Coastal Plain, within the Perth Metropolitan Region, 48% of the Quindalup Complex is estimated to remain, 20% of which is proposed for protection through Bush Forever (Government of Western Australia 2000a).

1.4 PROTECTION OF FLORA AND VEGETATION

Flora species are protected formally and informally at both the National and State level by various legislative and non-legislative measures, which are discussed below:

Legislative Protection

- *Environment Protection and Biodiversity Conservation Act 1999* (Cth): a Federal Act;
- *Wildlife Conservation Act 1950* (WA): a State Act; and
- *Environmental Protection Act 1986* (WA): a State Act.

Non-Legislative Protection

- Western Australian Department of Environment and Conservation (DEC) Priority lists for flora and vegetation; and
- informal recognition of locally significant populations

A short description of these measures is given below, and definitions of the species' conservation codes and ecological community categories they use, and those used by the DEC, are provided in Appendix A.

Environment Protection and Biodiversity Conservation Act 1999 (Cth)

The *Environment Protection and Biodiversity Conservation Act 1999* (the EPBC Act) aims to protect matters of national environmental significance, which are detailed in Appendix A.

Under the EPBC Act, the Commonwealth Department of Sustainability, Environment, Water, Populations and Communities (DSEWPaC) lists threatened species and Threatened Ecological Communities (TECs) in certain categories determined by criteria set out in the Act (www.environment.gov.au/epbc/index.html).

The Act provides for a national environmental assessment and approvals process for proposed actions likely to affect the prescribed matters of national environmental significance. If a proposed action is approved subject to certain conditions, the proponent of the action does not contravene the Act if the action is carried out in accordance with the conditions imposed.

Projects likely to cause impacts on matters of national environmental significance (as defined in the EPBC Act – see Appendix A) should be referred to DSEWPAC for assessment under the EPBC Act.

Wildlife Conservation Act 1950 (WA)

The Western Australian DEC recommends flora taxa for listing under the provisions of the *Wildlife Conservation Act 1950* (WC Act) as protected according to its need for protection (see Appendix A).

Flora species are given Declared Rare status when their populations are geographically restricted or are threatened by local processes. In addition, under the WC Act, by Notice in the Western Australian Government Gazette of 9 October 1987, all native flora (spermatophytes, pteridophytes, bryophytes and thallophytes) is protected throughout the State.

The Act makes it an offence to 'take' threatened species without an appropriate licence. There are financial penalties for contravening the Act.

Environmental Protection Act 1986 (WA)

Declared Rare Flora (DRF) and TECs are given special consideration in environmental impact assessment, and are Environmentally Sensitive Areas (ESAs) under the *Environmental Protection Act 1986* (EP Act) and the *Environmental Protection (Clearing of Native Vegetation) Regulations 2004*.

The protection of DRF and TECs is a 'clearing principle' for assessing applications for permits to clear native vegetation, where exemptions for a clearing permit under the *Environmental Protection (Clearing of Native Vegetation) Regulations 2004* do not apply. There are substantial penalties for unlawfully damaging ESAs.

DEC Priority Lists

The DEC lists 'Priority' flora species that have not been assigned statutory protection under the WC Act, but which are under consideration for declaration as 'Rare Flora' under the Act. Species assessed as Priorities 1-3 are in urgent need of further survey, whilst Priority 4 species require monitoring every 5-10 years (see Appendix A for definitions).

In addition, the DEC maintains a list of Priority Ecological Communities (PECs) which identifies those communities that need further investigation before possible nomination for TEC status. The DEC identifies and lists vegetation communities believed to be threatened. Once listed, a community is a PEC, but only when endorsed by the Minister for the Environment does it become a TEC, and

therefore becomes protected as an ESA under native vegetation clearing regulations (see Appendix A for definitions).

Informal Recognition of Flora and Vegetation

Certain populations or communities may be of local significance or interest because of their patterns of distribution and abundance. For example, flora may be locally significant because they are range extensions to the previously-known distribution or are newly-discovered taxa (and therefore have the potential to be of more than local significance). In addition, many species are in decline as a result of threatening processes (primarily land clearing), and relict populations of such species assume local importance.

Despite the lack of any formal protection for species in this category, project proponents are strongly advised to be aware of and to be sensitive to community concerns as to locally significant species or communities.

1.5 INTRODUCED SPECIES

The Environmental Weed Strategy for Western Australia (Department of Conservation and Land Management 1999) contains criteria for the assessment and ranking of weeds in terms of their environmental impact on biodiversity (Appendix B). The Strategy defines environmental weeds as 'plants that establish themselves in natural ecosystems and proceed to modify natural processes, usually adversely, resulting in the decline of the communities they invade.'

Plants may also be 'Declared' by the Agriculture Protection Board under the *Agriculture and Related Resources Protection Act 1979* (WA) (ARRP Act). Declared Plants are gazetted under five categories (P1-P5), which define the action required. Details of the definitions of these categories are provided in Appendix B. A declaration may apply to the whole State, to districts, individual properties or even to single paddocks. If a plant is Declared, landholders are obliged to control that plant on their properties (Department of Agriculture 2009).

1.6 BUSH FOREVER

Bush Forever is a State Government Policy and program which identifies 51,200 hectares of regionally significant bushland for protection, covering 26 vegetation complexes. This amounts to about 18% of the original vegetation on the Swan Coastal Plain portion of the Perth Metropolitan Region, and excludes local conservation reserves.

Regionally significant bushland has been identified on the basis of criteria relating to its conservation values. Important among these criteria is the achievement, where possible, of a comprehensive representation of all the ecological communities originally occurring in the region, principally through protecting a target of at least 10 per cent of each vegetation complex (Government of Western Australia, 2000a).

The Government of Western Australia has endorsed Bush Forever as the means of seeking the appropriate protection and management of areas of regionally significant bushland on the Swan Coastal Plain Portion of the Perth Metropolitan Region and a balance between environmental, social and economic objectives. As an endorsed government policy it is used as a basis for decision-making and an agreed framework for the protection and management of Bush Forever Sites through the implementation mechanisms identified in the plan (Government of Western Australia, 2000a).

The survey area is Bush Forever Site Number 355; the Point Peron and Adjacent Bushland, Peron/Shoalwater Bay.

2 METHODS

The aim of the assessment was to design and undertake the targeted spring vegetation and flora survey in the survey area to supplement previous biological survey work, in particular that undertaken by Bennett Environmental Consulting Pty Ltd (Bennett 2005).

All flora surveys undertaken by ENV are designed to be compliant with the Environmental Protection Authority (EPA) requirements for the environmental surveying and reporting of flora surveys in Western Australia, as set out in the following documents:

- *Terrestrial Biological Surveys as an Element of Biodiversity Protection. Position Statement No. 3* (EPA 2002); and
- *Guidance for the Assessment of Environmental Factors – Terrestrial Flora and Vegetation Surveys for Environmental Impact Assessment in Western Australia* (EPA 2004)

ENV then assesses and reports the results of its surveys with particular regard to the provisions of the EPBC Act, WC Act and EP Act.

The methodology for the work involved the following key steps:

PHASE 1 –SURVEY DESIGN

A vegetation and flora survey was completed within the survey area in August 2005 by Bennett Environmental Consulting Pty Ltd (Bennett). Prior to this a flora and vegetation survey was conducted by Keating and Trudgen in October 1986. Consequently a targeted spring survey was considered necessary to complete the botanical data for the survey area, with particular reference to determining the presence of Declared Rare and/or Priority Flora.

In addition, Bennett (2005) identified one potential Threatened Ecological Community (SCP30a *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands) at three sites within the survey area. Re-assessment of the species composition and condition of these sites is necessary to determine whether they represent viable examples of the TEC. This re-assessment includes re-visiting quadrats and recording the necessary data to confirm the floristic community type and to discuss its status as a TEC.

PHASE 2 - DESKTOP SURVEY

The purpose of a desktop survey is to obtain information on flora and vegetation constraints that may be present at the site. The tasks involved undertaking a desktop investigation on regional vegetation complexes, soils and landforms and

Bush Forever reference sites, and the review of available reports addressing the site.

In addition, a Geographic Information Systems (GIS) database search was submitted to the DEC to ascertain the locations of any DRF or Priority Flora species that have been recorded within the survey site and its surrounding areas. The search was within a 15km radius using coordinates from 32° 0' - 32° 25' S 115° 33' - 115° 51' E (GDA94).

A similar process is also undertaken to establish the locations of any additional TECs or Priority Ecological Communities (PECs) known from the area. Generally the search coordinates allow for a buffer of 15 km around the site.

Surveys undertaken by Keating and Trudgen (1986) and Bennett (2005) were also reviewed for flora species, vegetation community and vegetation condition information.

PHASE 3 – FIELD SURVEY

The field survey was undertaken in spring, between the 27th and 29th October 2009. The field survey consisted of:

- Establishment of two permanent quadrats within each of the inferred FCTs as described by Bennett (2005), at the same quadrat locations previously assessed. Rescoring of quadrat species data on two or more occasions is recommended by the DEC for performance of statistical analysis. Two quadrats within a vegetation community are recommended to adequately describe each community.
- Production of an inventory (list) of plant taxa associated with the potential TEC SCP30a. Each FCT has a list of common and indicator species which are diagnostic of the FCT ;
- searching for and mapping the location of any DRF and Priority Flora species and any other flora of local or taxonomic significance along 100 m spaced grid transects throughout the study area and intensified in their known habitats;
- providing a description and map/s of vegetation condition over the study area (as per the Keighery (1994) condition rating scale); and
- identifying and mapping any TECs within the study area.

PHASE 4 – DATA ANALYSIS AND REPORTING

Where field identification of plant taxa was not possible, specimens were collected systematically for later identification at the Western Australian

Herbarium (WAH) by comparison with the reference collection and use of identification keys.

Once all species were identified, plant taxa found within the study area were compiled into an inventory listed chronologically by family number. This list is checked against Florabase and the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) listing of Threatened species to determine species' conservation status and to determine records of species that constitute range extensions.

Where a significant species is found, an additional database search is submitted to the DEC to obtain data on the species' range and population details to determine the significance of the species at the site in a regional context.

Following completion of reporting, significant species will be vouchered with the Western Australian Herbarium (WAH) in accordance with conditions of DEC flora licences.

The data from the quadrats is compared against Gibson *et al.*'s (1994) Swan Coastal Plain (SCP) Floristic Community Types (FCTs) database using multivariate statistical analysis. This comprises transformation and normalisation of data and computation of a similarity matrix based on Bray-Curtis similarity to indicate similarity of the quadrat data (and therefore the vegetation unit) to documented FCTs.

The results of the statistical analysis are then considered in light of other site characteristics such as soils, landform and the presence of dominant, common or indicator species to draw conclusions on the FCT present.

When a conclusion has been reached over the FCT present, where this FCT is a TEC or PEC, further discussion of the site's characteristics, in particular vegetation condition and structure, size and edge to perimeter ratio are considered to determine whether the vegetation represents a viable example of the TEC or PEC.

2.1 FLORA SURVEY VARIABLES

It is important to note the specific variables imposed on individual surveys. Variables are often difficult to predict, as is the extent to which they influence survey effort. Survey constraints of the project area flora and vegetation survey are detailed in Table 1.

Table 1: Limitations and Constraints Associated with the project area Flora and Vegetation Survey

Variable	Impact on Survey Outcomes
Access Problems	No access problems were encountered during the field survey.
Experience levels	<p>The botanists who executed these surveys were practitioners suitably qualified in their respective fields.</p> <ul style="list-style-type: none"> • Coordinating Botanist: Narelle Whittington (Senior Botanist); • Field Staff: Narelle Whittington (Senior Botanist), Natalie Pawley (Botanist) and Peter Jobson (Taxonomist/Senior Botanist); • Taxonomy: Narelle Whittington (Senior Botanist); • Data Interpretation: Narelle Whittington (Senior Botanist);
Timing ¹ , weather, season.	<p>The survey was undertaken in Spring, between 27-29th October 2009. The area (Garden Island) had received 483.8 millimetres of rain in the year to date (January to October 2009; Bureau of Meteorology 2009).</p> <p>Flora composition changes over time, with flora species having specific growing periods, especially annuals and ephemerals (some plants lasting for a markedly brief time, some only a day or two). Therefore the results of future botanical surveys in this location may differ from the results of this survey.</p>
Completeness	<p>Species that were insufficiently mature or dead were identified in the field to genus or family level only (where possible).</p> <p>A comprehensive species list has not been prepared for areas that do not constitute a natural vegetation area, such as gardens or areas that have been totally cleared.</p>
Determination	<p>This survey makes inferences about vegetation types that have the potential to be TECs. However, a decision as to the presence or absence of TECs at the site remains the responsibility of the DEC's Species & Communities Branch.</p> <p>The taxonomy and conservation status of the Western Australian flora are dynamic. This report was prepared in reliance on taxonomy and conservation current at the time issuing, but it should be noted this may change.</p>

¹ EPA Guidance Statement 51 (2004) stipulates that flora and vegetation surveys should be undertaken following the season that contributes the greatest rainfall in the region. In the South-west Province the main rain is in winter, requiring surveys to be undertaken in spring. Short-term variations in normal weather patterns (e.g. drought) may necessitate supplementary survey work at other times of year or in later years to take into account temporal changes in diversity.

2.2 PERMITS

Specimens collected during the survey were taken by permit of and subject to the conditions of the following licences issued under sections 23C and 23F of the Wildlife Conservation Act:

- SL008739 and 72-0910 Narelle Whittington;
- SL008486 and 46-0910 Natalie Pawley; and
- SL008532 Peter Jobson.

3 RESULTS

3.1 DATABASE SEARCH

A database search of the area resulted in four Declared Rare and 15 Priority Flora species being identified as potentially occurring in the area. For a comprehensive list of species found during the database search, please refer to Appendix C.

The database search determined that four TECs are known to occur in the area (see Appendix C).

3.2 FIELD SURVEY - FLORA

3.2.1 Flora

The Bennett (2005) survey recorded an additional 21 taxa, which were not recorded by ENV. These additional species were recorded within quadrats that were not rescored by ENV. The majority of these species are weeds (15 species). These species were not used in the statistical analysis and 19 have not been discussed below. Two of these species are significant flora and have been discussed (*Allocasuarina lehmanniana* and *Hibbertia cuneiformis*). All of these species are listed in Appendix D.

Seventy five taxa comprising 37 families and 65 genera were recorded in the survey area (41 native flora taxa and 34 introduced). One species was unable to be identified below family level while one species was unable to be identified past genus level. Refer to Appendix D for the flora species matrix and Appendix E for the flora survey field datasheets and site photographs.

The plant families most frequently recorded from the survey were as follows:

- Poaceae 13 species;
- Fabaceae 10 species;
- Asteraceae six species; and
- Myrtaceae five species

The plant genera most frequently recorded from the survey were as follows:

- *Acacia* five species; and
- *Euphorbia* three species.

The average species richness is 15 taxa per quadrat.

3.2.2 Protected Flora

No Threatened species pursuant to the EPBC Act were located during the survey.

No plant taxa gazetted as Declared Rare pursuant to the WC Act were located in the survey area.

No Priority Flora species were located in the survey area.

3.2.3 Locally Significant Flora

Bush Forever lists species that are considered to be of particular interest in the Perth Metropolitan Area. Other than DRF or Priority Flora these species may be of interest due to being restricted in distribution, endemic to a particular location or have some other distinctive feature (Government of Western Australia 2000b) and are presented as being locally significant flora.

Species that were located by ENV and Bennett Consulting (2005) within the Cape Peron Survey area that are considered to be of significance and the reason for that significance as per Bush Forever are listed in Table 2. See Figure 4 for significant flora locations.

Table 2: Significant Flora Locations

Species	Significance Category	Location
<i>Agonis flexuosa</i> var. <i>flexuosa</i>	At northern extension of Known range; Significant population.	ENV 378 408E 6427831N 376977E 6428639N
<i>Allocasuarina lehmanniana</i>	Significant population.	ENV did not record this species Bennett recorded this species but its co-ordinates were not recorded
<i>Callitris preissii</i>	Significant population. Taxa endemic to the Swan Coastal Plain in the Perth Metropolitan Region.	ENV 387408E 6427831N 376415E 6428720N
<i>Diplolaena dampieri</i>	At northern extension of known range; significant population.	ENV 376477E 642883N Bennett 376477E 6428883N 376894E 6428773N
<i>Hibbertia cuneiformis</i>	At northern extension of known range; significant	ENV did not record this species

Species	Significance Category	Location
	population.	Bennett 377968E 6426050N
<i>Melaleuca lanceolata</i>	Disjunct population; significant population.	ENV 376890E 6428951N 376416E 6428721N Bennett 376930E 6428962N

3.2.4 Flora Potentially Sensitive to Groundwater Changes

The species listed in Table 3 are those that are susceptible to either changes in the level of the water table, or to changes in the water quality within the water table.

Table 3: Flora Potentially Sensitive to Groundwater Changes

Family	Species	Susceptible to Changes in Groundwater Levels	Susceptible to Changes in Groundwater Quality
Cupressaceae	<i>Callitris preissii</i>		X
Poaceae	<i>Spinifex hirsutus</i>	X	
Poaceae	<i>Spinifex longifolius</i>	X	
Cyperaceae	<i>Ficinia nodosa</i>	X	
Cyperaceae	<i>Lepidosperma gladiatum</i>	X	
Cyperaceae	<i>Lepidosperma</i> sp. Coastal Dune (R. J. Cranfield 9963)	X	
Restionaceae	<i>Desmocladus flexuosus</i>		X
Myrtaceae	<i>Agonis flexuosa</i>	X	
Myrtaceae	<i>Eucalyptus gomphocephala</i>	X	
Myrtaceae	<i>Melaleuca lanceolata</i>	X	
Frankeniaceae	<i>Frankenia pauciflora</i>	X	
Epacridaceae	<i>Leucopogon parviflorus</i>		X

3.2.5 Introduced Flora

The table below (table 4) contains the weed species identified during the field survey, with their ratings and criteria according to the Environmental Weed Strategy for Western Australia (refer to Appendix B for the criteria used for ranking).

Table 4: Weed Species Identified

Taxon	Common Name	Criteria			
		Rating	Invasiveness	Distribution	Impacts
<i>*Anagallis arvensis</i>	Pimpernel	Moderate	Yes	Yes	
<i>*Arctotheca calendula</i>	Cape Weed	Moderate	Yes	Yes	
<i>*Asparagus asparagoides</i>	Bridal Creeper	High	Yes	Yes	Yes
<i>*Avena barbata</i>	Bearded Oat	Moderate	Yes	Yes	
<i>*Bromus diandrus</i>	Great Brome	High	Yes	Yes	Yes
<i>*Cakile maritima</i>	Sea Rocket	Moderate	Yes	Yes	
<i>*Carpobrotus edulis</i>	Hottentot Fig	Moderate	Yes	Yes	
<i>*Crassula glomerata</i>		Moderate	Yes	Yes	
<i>*Cuscuta epithymum</i>	Lesser Dodder	Moderate	Yes	Yes	
<i>*Cynodon dactylon</i>	Couch	Moderate	Yes	Yes	
<i>*Ehrharta calycina</i>	Perennial Veldt Grass	High	Yes	Yes	Yes
<i>*Ehrharta longiflora</i>	Annual Veldt Grass	Moderate	Yes	Yes	
<i>*Eragrostis curvula</i>	African Love Grass	High	Yes	Yes	Yes
<i>*Euphorbia paralias</i>	Sea Spurge	Moderate	Yes	Yes	
<i>*Euphorbia peplus</i>	Petty Spurge	Moderate	Yes	Yes	
<i>*Euphorbia terracina</i>	Geraldton Carnation Weed	High	Yes	Yes	Yes
<i>*Foeniculum vulgare</i>	Fennel	N/A	N/A	N/A	N/A
<i>*Fumaria capreolata</i>	Whiteflower Fumitory	Mild			Yes
<i>*Geranium molle</i>	Dove's Foot	Low	N/A	N/A	N/A

Taxon	Common Name	Criteria			
		Rating	Invasiveness	Distribution	Impacts
	Cranesbill				
<i>*Hyparrhenia hirta</i>	Tambookie Grass	High	Yes	Yes	Yes
<i>*Lagurus ovatus</i>	Hare's Tail Grass	High	Yes	Yes	Yes
<i>*Lolium rigidum</i>	Wimmera Ryegrass	Moderate	Yes	Yes	
<i>*Melilotus indicus</i>		N/A	N/A	N/A	N/A
<i>*Olea europaea</i>	Olive	Moderate	Yes	Yes	
<i>*Pelargonium capitatum</i>	Rose Palargonium	High	Yes	Yes	Yes
<i>*Pennisetum clandestinum</i>	Kikuyu Grass	Moderate	Yes	Yes	
<i>*Plantago lanceolata</i>	Ribwort Plantain	Low	N/A	N/A	N/A
<i>*Rhamnus alaternus</i>	Buckthorn	Moderate	Yes	Yes	
<i>*Romulea rosea</i>	Guildford Grass	High	Yes	Yes	Yes
<i>*Schinus terebinthifolius</i>		N/A	N/A	N/A	N/A
<i>*Sonchus oleraceus</i>	Common Sowthistle	Moderate	Yes	Yes	
<i>*Tetragonia decumbens</i>	Sea Spinach	Moderate	Yes	Yes	
<i>*Trachyandra divaricata</i>	Onion Weed	Mild		Yes	
<i>*Urospermum picroides</i>	False Hawkbit	Moderate	Yes	Yes	

One Declared Plant species, **Asparagus asparagoides*, was found in the study area. This species is listed as Priority 1 for the whole State.

3.3 FIELD SURVEY – VEGETATION

3.3.1 Floristic Community Types

The Quindalup Complex on the Quindalup landform is recognised as being composed of 12 SCP and supplementary FCTs. These are:

SCP 17: *Melaleuca raphiophylla* – *Gahnia trifida* seasonal wetlands;

SCP 19: Sedgelands in Holocene dune swales;

- SCP 29a:** Coastal shrublands on shallow sands;
- SCP 29b:** *Acacia* shrublands on taller dunes;
- SCP 30a** *Callitris preissii* and/or *Melaleuca lanceolata* forests and woodlands;
- SCP 30c** Woodlands and shrublands on Holocene dunes (re-allocated from 30c);
- SCP 30b:** Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands;
- SCP S11:** Northern *Acacia rostellifera* – *Melaleuca acerosa* shrublands;
- SCP S12:** Rottnest Island *Melaleuca lanceolata* and/or *Callitris preissii* forests and woodlands;
- SCP S13:** Northern *Olearis axillaris* – *Scaevola crassifolia* shrublands;
- SCP S14:** *Spinifex longifolius* grassland and low shrublands;
- SCP S 15:** Weed group. Not allied with any supergroup.

The vegetation units of the site have been mapped previously (Bennett 2005) and these have been inferred to eight Floristic Community Types (FCTs). The eight inferred Floristic Community types are as follows:

- FCT SCP16:** Highly saline seasonal wetlands
- FCT SCP29a:** Coastal shrublands on shallow sands
- FCT SCP29b:** *Acacia* shrublands on taller dunes
- FCT S13:** Northern *Olearia axillaris* – *Scaevola crassifolia* shrublands
- FCT S14:** *Spinifex longifolius* grasslands and low shrublands
- FCT SCP30a:** *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands
- FCT SCP30b:** Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands
- FCT S15:** Weed Group

ENV established sixteen quadrats within the eight FCTs within the survey site. To ensure the survey is undertaken in accordance with Guidance Statement 51 (EPA 2004), replication quadrats were established to increase the number of

species for each vegetation unit. Opportunistic collections were also recorded for each of the units to add to the species total.

To conform that these eight FCTs occur within the study area, data analysis was undertaken.

Determination of Floristic Community Type by Similarity

The analysis suggested that the site appeared to belong to several FCTs. The results of the data analysis are shown in Table 5. The data analysis results illustrating the Bray-Curtis similarity is illustrated within the dendrogram included as Appendix G.

Table 5: Summary of PRIMER Analysis

Floristic Community Type (Quadrat)	Analysis FCT	% Similarity
SCP 29b (Q1 & 8)	30a	33
SCP 16 (Q3 & 5)	29a	24
SCP 30a (Q2, 11 & 12)	30a	38
S13 (Q6 & 15)	30a	24
S15 (Q7)	19	18
SCP 30b (Q9 & 10)	30a	19
SCP 29a (Q13 & 16)	30a	30
S14 (Q4 & 14)	29a	27

The results suggest that the FCTs that have been previously inferred vary from the data analysis. Importantly, all but one site have low similarity percentages, with them being below 30%, which means that further investigation into the quadrats' characteristics are needed in order to allocate the vegetation to FCTs.

It was found, however, that all the sites are clearly related to Quindalup Complex communities.

Due to the inconclusive results of the statistical analysis there was a need to further analyse the data to clarify what FCTs best correlate with the quadrats independently of the statistical analysis. This involves reviewing site data for other factors that are diagnostic for FCTs, including the presence of indicator species, soil types and landform position.

Table 6: Floristic Community Type Determination

Vegetation Unit (Quadrat)	Primer Analysis	% Similarity	Comments	ENV inferred	% Similarity
SCP 29b (Q1 & 8)	30a	33	The main indicator species for SCP 30a is absent from the sites and is more characteristic of FCT 29b	29b	31
SCP 16 (Q3 & 5)	16 and 29a	24	ENV concurs with the analysis results that it is characteristic of both SCP 16 and 29a.	16 and 29a	24
SCP 30a (Q2, 11 & 12)	30a	38	ENV concurs with the analysis results	30a	38
S13 (Q6 & 15)	30a	24	The main indicator species for SCP 30a is absent from the sites and even though the data for the supplementary community types is not available for analysis, the site is characteristic of S13	S13	Not available
S15 (Q7)	19	18	ENV disagrees with the prospect of it being SCP 19 as the vegetation unit lacks the characteristics and dominant species of this FCT. The site is dominated by weeds with minimal native species making it more affiliated with S15	S15	Not available

Vegetation Unit (Quadrat)	Primer Analysis	% Similarity	Comments	ENV inferred	% Similarity
SCP 30b (Q9 & 10)	30a	19	The main indicator species for SCP 30a is absent from the sites however the dominant indicator species of SCP 30b is present and the quadrats occur on the correct landform.	30b	-
SCP 29a (Q13 & 16)	30a	30	ENV disagrees with the prospect of it being SCP 19 as the vegetation unit lacks the characteristics and dominant species of this FCT. The site have more affiliation with 29a based on species present.	29a	28
S14 (Q4 & 14)	29a	27	Due to the dominant species present the two quadrats are not compatible with the analysis results and are more characteristic of FCT S14.	S14	Not available

See Figures 4 and 5 for Quadrat locations and FCT locations.

3.3.2 Priority Ecological Communities and Threatened Ecological Communities

FCT SCP30a *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands is listed as a TEC and is listed as Vulnerable by the State

FCT SCP30b Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands is listed as a Priority Ecological Community (Priority 3).

FCT SCP29b *Acacia* shrublands on taller dunes is listed as a Priority Ecological Community (Priority 3).

3.3.3 Vegetation Condition

The condition scale commonly used in the Perth metropolitan area and Bush Forever (Government of Western Australia 2000), was used for this assessment. The definition of the condition scales is in Appendix F.

The condition of the vegetation on site varies between Very Good and Completely Degraded (excluding development areas). Refer to Figure 6 for a map of bushland condition.

The high variability of the condition is reflected by the fragmentation of the area by different infrastructure, roads, tracks, weeds and rubbish. The volume of people that use the area everyday has contributed to the degradation of the vegetation both directly through trampling and spread of weeds and indirectly through the need for additional infrastructure such as roads and amenities. In the majority of cases the vegetation is in Good condition however without protection or active remediation the condition of the site will further degrade.

3.3.4 Bush Forever

The entire survey site is mapped as a Bush Forever site; the Point Peron and Adjacent Bushland, Peron/Shoalwater Bay, Site Number 355. The Bush Forever site includes all the remnant vegetation within the survey site equating to approximately 107.1 ha.

Government of Western Australia (2000b) states a detailed survey was undertaken of the site by Keating and Trudgen in 1986, which resulted in 60% of the flora taxa being sampled with no significant species being found.

The site meets six specific coastal reserve criteria, these are:

- Quindalup Dune types: youngest, older and beach ridge plain
- Continuing natural processes: 174.5 ha (107.1 ha of bushland) of Quindalup Dunes extending to 3.1 km inland from the point
- Shoreline: soft (sandy) and Hard (rock)
- Linkage: contains Quindalup/Spearwood Dunes (Tamala Limestone) interface; roads and developments fragment site
- Vegetation: typical Quindalup/Spearwood units
- Habitats: significant reptile species

4 DISCUSSION

Flora

During the survey, a total of 75 taxa, from 37 families and 65 genera were recorded within the survey area (41 native flora taxa and 34 introduced taxa). Of these, no Threatened species pursuant to the EPBC Act, Declared Rare Flora pursuant to the WC Act or Priority Flora species were located.

ENV considers the number of flora taxa reported here is not an accurate representation of the potential amount of flora species present within the survey site. ENV only established 16 quadrats within FCTs and did not survey vegetation units which would have created a more robust species list and thus a better representation of the flora species present.

Six locally significant species were found across the site (details in Section 3.2.3). The reasons for their significance are attributed to them occurring at the northern extension of their known range and are considered a significant population according to Government of Western Australia (2000b). These species are not listed for protection, however are considered to be of interest.

A number of species have also been identified as potentially sensitive to changes in groundwater levels and or quality. Plants such as *Ficinia nodosa* and *Frankenia pauciflora* commonly occur in areas that are seasonally inundated, requiring their roots to be in waterlogged conditions for short periods. The lowering of the water table from current levels could reduce the regularity or occurrence of these low-lying areas experiencing inundation. Community composition could change as a result.

Larger tree species such as *Agonis flexuosa*, *Eucalyptus gomphocephala* and *Melaleuca lanceolata* have shallow root systems and are commonly found in low lying areas, with raised water table levels. The lowering of the water table could induce stress and potentially cause the death of individuals of these plants.

Threatened, Declared Rare and Priority Flora

No Threatened species pursuant to the EPBC Act, Declared Rare Flora pursuant to the WC Act or Priority Flora species were located in the survey area during the survey. Consequently, based on the legislative framework there are no species of international, National or State significance located within the project area.

ENV considers that the potential for the site to contain Declared Rare or Priority Flora is low. This is because the survey was undertaken in spring at peak flowering time and due to the nature of the project area the entire site was able to

be extensively traversed on foot. If any Threatened, Declared Rare or Priority flora species were present on site they would have been easily located.

Dodonaea hackettiana, which is the only priority flora that has been previously recorded within the project area was not recorded by ENV at the time of the survey. The species was also not located by Bennett Consulting during the 2005 survey. Considering the species was not found in the location that it was previously known to occur, nor the rest of the Cape Peron survey area by two intensive surveys undertaken by ENV and Bennett Consulting, it can be assumed that it was misidentified or the individuals have subsequently died.

Weeds

Thirty four weed species were identified during the survey. Had a extensive weed survey been undertaken on the Degraded to Completely Degraded areas then it is probable the number of species present would have been higher. Weed species were only recorded if they were dominant, occurred within a quadrat, or are listed as a Declared Plant. Due to the survey site being used heavily by people, weeds are the main disturbance factor of the native vegetation (besides clearing for infrastructure). The weeds present are typical for urban sites.

One Declared Plant species, **Asparagus asparagoides*, listed by the ARRP Act was found in the study area. This species is listed as Priority 1 for the whole State.

Vegetation Complexes

The Environmental Protection Authority (EPA) recognises that native vegetation complexes which have less than 10% of pre-European clearing extent remaining in the Bush Forever study area within the Perth Metropolitan Region (PMR) may be considered regionally significant (EPA 2006). Proposals that would impact on a vegetation complex with 10% or less remaining may be formally assessed by the EPA (EPA 2006).

The site is mapped as Quindalup Complex: Coastal dune complex consisting mainly of two alliances – the standard fore-dune alliance and the mobile and stable dune alliance. Local variations include the low closed forest of *Melaleuca lanceolata* – *Callitris preissii* and the closed scrub of *Acacia rostellifera*.

Bush Forever gives an estimate of the percentage of each complex that remains within the Bush Forever study area compared to its pre-European settlement extent, so an estimate of the scarcity of each complex can be determined. On the Swan Coastal Plain, within the PMR, 48% of the Quindalup Complex is estimated to remain, 20% of which is proposed for protection through Bush Forever which aims to retain at least of 10% of each vegetation complex within the Bush Forever policy area (Government of Western Australia 2000a).

As the entire survey site is included in the Bush Forever site, the remnant vegetation (equating to approximately 107.1 ha) is classified as regionally significant and contributes to the retention targets documented in Bush Forever (Government of Western Australia 2000a).

The site does have some existing protection and it is recommended in the Bush Forever report (Government of Western Australia 2000b) that the care, control and management of the site for conservation purposes within the Rockingham Lakes Regional Park is endorsed. Part of the site is designated as government lands (including existing and proposed public utilities).

Floristic Community Types

The vegetation identified in this assessment can broadly be related to eight of Gibson *et al* (1994) Floristic Community Types, as described in Section 3.3.1.

Due to the inconclusive results of the statistical analysis, attributed to low species diversity, the limited survey work that has been undertaken on the Quindalup Dune System and the condition of the vegetation in some instances, there was a need to review the species data and site information to clarify what FCTs best correlate with the quadrats in collaboration with the data analysis.

The following FCTs have been identified by this assessment as occurring on site:

FCT SCP16:	Highly saline seasonal wetlands
FCT SCP29a:	Coastal shrublands on shallow sands
FCT SCP29b:	<i>Acacia</i> shrublands on taller dunes
FCT S13:	Northern <i>Olearia axillaris</i> – <i>Scaevola crassifolia</i> shrublands
FCT S14:	<i>Spinifex longifolius</i> grasslands and low shrublands
FCT SCP30a:	<i>Callitris preissii</i> (or <i>Melaleuca lanceolata</i>) forest and woodlands
FCT SCP30b:	Quindalup <i>Eucalyptus gomphocephala</i> and/or <i>Agonis flexuosa</i> woodlands
FCT S15:	Weed Group

One TEC, SCP30a *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands was identified as occurring within the survey area at three locations. This FCT is listed as Vulnerable by the State and is not listed by the Commonwealth. This FCT is represented by quadrats two, eleven and twelve. This TEC is considered to be of State significance.

Three quadrats were established within the SCP30a to record the viability of the vegetation community at those locations. The SCP30a located on the boundary of the school has been mapped as being in Good condition, however there are important factors to take into account that contribute to the vegetation community's survival. The site is very small (7 m x 30 m) and is located between the coastal sand dunes and a school oval and there is a pathway that dissects the vegetation to allow students to access the beach. There is no native understorey present and since the 2005 survey all the *Callitris preissii* have died. This leads to the conclusion that the vegetation community is subject to recent degrading factors. ENV concludes that this particular example is not a viable representation of this FCT and does not represent the TEC.

The site surrounding the small grassed area on the corner of Memorial Drive and Safety Bay Road is in Good to Degraded condition and is partly located within the impact area (Figure 7). The TEC is approximately 1.5ha in size, of which 0.4 ha is in degraded condition and 1.1 ha is in Good condition. Approximately 709 m² of TEC in Good condition occurs within the impact area, this represents approximately 6% of the 1.1 ha of TEC in Good condition.

Due to its location and being surrounded by native vegetation, the site has the potential to be remediated to improve its condition. There are many informal tracks that dissect the area and rubbish has been dumped adjacent to the site so these degrading influences will need to be resolved if the vegetation community is to be conserved.

The third site located at the base of the western-most car park is surrounded by dunal vegetation and therefore the degrading factors are at a minimum, thus the site is a viable representation of the vegetation community. However there is a walking track that runs adjacent to the site and therefore issues such as people deviating off the path and the spread of weeds needs to be resolved to protect the community.

Two of the FCTs are listed as Priority 3 PECs by the State, FCT SCP30b Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands (quadrats 9 and 10; Figure 4). The occurrences of this FCT are in Good condition but are under threat by direct human disturbances (trampling, rubbish and weeds) and therefore would benefit from fencing and weed management. FCT SCP29b *Acacia* shrublands on taller dunes (quadrats 1 and 8; Figure 4) occurs across the majority of the site. Portion of this FCT are in Very Good condition and efforts should be made to preserve these in their current state. As PECs these areas of vegetation are considered to be of a Regional to State level of significance.

Vegetation Condition

The condition of the vegetation on site varies from Very Good to Completely Degraded, with the majority of the site in Good condition. There is no vegetation within the project area that is considered in excellent condition and there are only patches of vegetation mapped as Very Good condition (see Figure 6). This can be attributed to the various land uses and the high human traffic throughout the area.

5 SUMMARY AND RECOMMENDATIONS

The flora and vegetation survey undertaken by ENV.Australia determined that:

- A total of 75 taxa, from 37 families and 65 genera were recorded within the project area (41 native flora taxa and 34 introduced taxa).
- No Threatened species pursuant to the EPBC Act, Declared Rare Flora pursuant to the WC Act or Priority Flora species were located. Consequently, based on the legislative framework there are no species of international, National or State significance located within the survey area.
- Six species identified as being locally significant occur within the survey area.
- Twelve species are identified as being flora potentially sensitive to changes in groundwater levels and/or quality.
- One Declared Plant species (**Asparagus asparagoides*) was found in the survey area:
- The survey area is mapped as a Bush Forever site and the vegetation within the survey area is therefore classified as being of regionally significant. The site is mapped as Quindalup Complex: Coastal dune complex, which exceeds the 10% recommended retention status for Western Australia by the EPA's *Position Statement No. 2*. Bush Forever recommends 20% of this complex for retention within the Perth Metropolitan area.
- The following FCTs have been identified as occurring on site:
 - **SCP16:** Highly saline seasonal wetlands
 - **SCP29a:** Coastal shrublands on shallow sands
 - **SCP29b:** *Acacia* shrublands on taller dunes
 - **S13:** Northern *Olearia axillaris* – *Scaevola crassifolia* shrublands
 - **S14:** *Spinifex longifolius* grasslands and low shrublands
 - **SCP30a:** *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands
 - **SCP30b:** Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands
 - **FCT S15:** Weed Group

- One TEC, FCT SCP30a *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands was identified as occurring on site. This FCT is considered to be of State significance.
- Two of the FCTs are listed as Priority 3 PECs; FCT SCP30b Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands and FCT SCP29b *Acacia* shrublands on taller dunes. These FCTs are of Regional to State significance.

5.1 RECOMMENDATIONS

ENV makes the following recommendations:

- The FCT located on the boundary of the school is not a good representation of the TEC and is not considered viable.
- The TEC site surrounding the small grassed area on the corner of Memorial Drive and Safety Bay Road is surrounded by native vegetation and therefore has the potential to be remediated to improve its condition. There are many informal tracks that dissect the area and rubbish has been dumped adjacent to the site so these degrading influences will need to be resolved if the vegetation community is to be conserved.
- The third TEC site located at the base of the western-most car park has a track that runs adjacent to the site and therefore issues regarding people deviating off the path and the spread of weeds needs to be resolved to protect the vegetation community.
- Two of the FCTs are listed as Priority 3 PECs by the state, FCT SCP30b Quindalup *Eucalyptus gomphocephala* and/or *Agonis flexuosa* woodlands and FCT SCP29b *Acacia* shrublands on taller dunes. PECs are not protected under legislation however should be treated as significant and taken into consideration during the planning phase.

6 REFERENCES

- Beard J S (1978) *Vegetation Survey of Western Australia – Pinjarra Sheet 1:250000*. Vegmap Publications, Perth, Western Australia.
- Beard J S (1990) *Plant Life of Western Australia*. Kangaroo Press.
- Bennett Environmental Consulting Pty Ltd (2005). Flora and Vegetation Point Peron Western Australia. Report prepared for Strategen Environmental Consultants.
- Bureau of Meteorology [BOM] (2009) *Daily Weather Observations*, Commonwealth of Australia. Online: www.bom.gov.au/climate
- Churchward H M and McArthur W M (1978) *Darling System, Landforms and Soils*. Division of Land Resources Management, CSIRO, Perth, Western Australia.
- Commonwealth of Australia (2001) *National Objectives and Targets for Biodiversity Conservation 2001 – 2005*. Department of Environment and Heritage.
- Department of Agriculture and Food Western Australia [DAFWA] (2008) *Declared Plants*. Online: www.agric.wa.gov.au [2009].
- Department of Conservation and Environment [DEC] (1980) *Atlas of Natural Resources Darling System Western Australia: Explanatory Text*. Department of Conservation and Environment, Western Australia.
- Department of Conservation and Land Management (1999) *Environmental Weed Strategy for Western Australia*. Department of Conservation and Land Management, Perth.
- Environmental Protection Authority [EPA] (2000) *Environmental Protection of Native Vegetation in Western Australia: Clearing of Native Vegetation with Particular Reference to Agricultural Areas, Position Statement No. 2*. EPA, Perth, Western Australia.
- Environmental Protection Authority [EPA] (2002) *Terrestrial Biological Surveys as an Element of Biodiversity Protection, Position Statement No. 3*. Environmental Protection Authority, Perth, Western Australia.
- Environmental Protection Authority [EPA] (2004) *Terrestrial Flora and Vegetation Surveys for Environmental Impact Assessment in Western Australia, Guidance Statement No. 51*. Environmental Protection Authority, Perth, Western Australia.

Environmental Protection Authority [EPA] (2006) *Level of Assessment for Proposals Affecting Natural Areas within the System 6 Region and Swan Coastal Plain Portion of the System 1 Region, Guidance Statement No. 10*. Environmental Protection Authority, Western Australia.

Gibson N, Keighery B, Keighery G, Burbidge A and Lyons M (1994) *A Floristic Survey of the Southern Swan Coastal Plain*. Prepared by Department of Conservation and Land Management and the Western Australian Conservation Council for the Australian Heritage Commission, Perth.

Government of Western Australia (2000a) *Bush Forever Volume 1: Policies, Principles and Processes*. Department of Environmental Protection, Perth, Western Australia.


Government of Western Australia (2000b) *Bush Forever Volume 2: Directory of Bush Forever Sites*. Department of Environmental Protection, Perth, Western Australia.


Hedde E M, Loneragan O W, and Havel J J (1978) *Vegetation Complexes of the Darling System (W.A.) – Pinjarra Sheet 1:250000*. Department of Conservation and Environment, Perth, Western Australia.

Keating, C. and Trudgen, M. (1986). A Flora and Vegetation Survey of the Point Peron – Lake Richmond Area. Unpublished Report for the State Planning Commission, Perth.

Western Australian Herbarium [WAH] (2009) *FloraBase - Information on the Western Australian Flora*. Department of Environment and Conservation, Perth. Online: <http://florabase.calm.wa.gov.au> [2010].

FIGURES



	Client: STRATEGEN	SITE LOCATION	Date: 3 February 2010
	Project: FLORA AND VEGETATION SURVEY OF THE MANGLES BAY AREA CAPE PERON		Scale: 1:1 Million
			Author: N.W. / S.C.
			Figure No. 1
			Plan No. CP-001



Author: N.Whittington
Drawn: S.Coleman
Status:
Job Number: 09.232

Client: **STRATEGEN**
Project: **FLORA AND VEGETATION SURVEY
OF THE MANGLES BAY AREA CAPE PERON**

FLORA SURVEY AREA

Date: 19 February 2010
Scale: 1:12 000
Figure No. **2**
Plan No. **CP-002**

A3



Legend

- Flora Quadrat
- Significant Flora



Author: N.Whittington	Client: STRATEGEN
Drawn: S.Coleman	Project: FLORA AND VEGETATION SURVEY OF THE MANGLES BAY AREA CAPE PERON
Status:	
Job Number: 09.232	

**QUADRAT AND
SIGNIFICANT FLORA LOCATIONS**

Date: 8 February 2010
Scale: 1:12 000
Figure No. 4
Plan No. CP-014



Floristic Community Types

- SCP16 - Highly saline seasonal wetlands.
- SCP29a - Coastal shrublands on shallow sands.
- SCP29b - (P3 PEC) *Acacia* shrublands on taller dunes.
- S13 - Northern *Olearia axillaris* - *Scaevola crassifolia* shrublands.
- S14 - *Spinifex longifolius* grasslands and low shrublands.
- SCP30a (TEC) - *Callitris preissii* (or *Melaleuca lanceolata*) forest and woodlands.
- SCP30b (P3 PEC) - Quindalup *Eucalyptus gomphocephala* and / or *Agonis flexuosa* woodlands.
- S15 - Weed Group.
- Development




Author: N.Whittington
Drawn: S.Coleman
Status:
Job Number: 09.232

Client: STRATEGEN
Project: FLORA AND VEGETATION SURVEY
OF THE MANGLES BAY AREA CAPE PERON

FLORISTIC COMMUNITY TYPE MAPPING

Date: 8 February 2010
Scale: 1:12 000
Figure No. 5
Plan No. CP-012



Vegetation Condition

- Very Good
- Very Good to Good
- Good
- Good to Degraded
- Degraded
- Degraded to Completely Degraded
- Completely Degraded



Author: N.Whittington
 Drawn: S.Coleman
 Status:
 Job Number: 09.232

Client: **STRATEGEN**
 Project: **FLORA AND VEGETATION SURVEY
 OF THE MANGLES BAY AREA CAPE PERON**

BUSHLAND AND CONDITION MAP

Date: 8 February 2010
 Scale: 1:12 000
 Figure No. **6**
 Plan No. **CP-013**

A3

Vegetation Condition

- Degraded SPC30a
- SCP30a (TEC)


SCALE 1 : 2 000

0 100

Metres

MAP GRID OF AUSTRALIA

Grid based on Transverse Mercator Projection
GDA 1994 MGA Zone 50



Client: **STRATEGEN**

Project: **FLORA AND
VEGETATION SURVEY
OF THE MANGLES BAY AREA
CAPE PERON**

AREA AND CONDITION OF TEC WITHIN THE IMPACT AREA

09.232

A4

Date: 5 March 2010

Scale: 1:2 000

Author: T.G. / S.C.

Figure No. **7**

Plan No. **CP-015**

APPENDIX A

DEFINITIONS OF DECLARED RARE / PRIORITY / THREATENED FLORA AND THREATENED / PRIORITY ECOLOGICAL COMMUNITIES

APPENDIX A

DEFINITIONS OF DECLARED RARE / PRIORITY / THREATENED FLORA AND THREATENED / PRIORITY ECOLOGICAL COMMUNITIES

A1: Categories of Declared Rare and Priority Flora

Conservation Code	Category
X	Declared Rare Flora - Presumed Extinct Taxa Taxa which have not been collected, or otherwise verified, over the past 50 years despite thorough searching, or of which all known wild populations have been destroyed more recently, and have been gazetted as such.
R	Declared Rare Flora - Extant Taxa "Taxa which have been adequately searched for and are deemed to be in the wild either rare, in danger of extinction, or otherwise in need of special protection and have been gazetted as such"
P1	Priority One - Poorly Known Taxa "Taxa which are known from one or a few (generally <5) populations which are under threat, either due to small population size, or being on lands under immediate threat, e.g. road verges, urban areas, farmland, active mineral leases, etc., or the plants are under threat, e.g. from disease, grazing by feral animals, etc. May include taxa with threatened populations on protected lands. Such taxa are under consideration for declaration as 'rare flora', but urgently need further survey."
P2	Priority Two - Poorly Known Taxa "Taxa which are known from one or a few (generally <5) populations, at least some of which are not believed to be under immediate threat (i.e. not currently endangered). Such taxa are under consideration for declaration as 'rare flora', but urgently need further survey."
P3	Priority Three - Poorly Known Taxa "Taxa which are known from several populations, and the taxa are not believed to be under immediate threat (i.e. not currently endangered), or known populations being large, and either widespread or protected. Such taxa are under consideration for declaration as 'rare flora' but need further survey."
P4	Priority Four - Rare Taxa "Taxa which are considered to have been adequately surveyed and which, whilst being rare (in Australia), are not currently threatened by any identifiable factors. These taxa require monitoring every 5-10 years."

Source: Department of Environment and Conservation (2009). *Western Australian Flora Conservation Codes*.

Department of Environment and Conservation, Perth, Western Australia. Online: <http://florabase.calm.wa.gov.au>.

A2: Categories of Threatened Flora Species

Category Code	Category
Ex	Extinct Taxa which at a particular time if, at the time, there is no reasonable doubt that the last member of the species has died.
ExW	Extinct in the Wild Taxa which is known only to survive in cultivation, in captivity or as a naturalised population well outside its past range; or it has not been recorded in its known and/or expected habitat, at appropriate seasons, anywhere in its past range, despite exhaustive surveys over a time frame appropriate to its life cycle and form.
CE	Critically Endangered Taxa which at a particular time, it is facing an extremely high risk of extinction in the wild in the immediate future, as determined in accordance with the prescribed criteria.
E	Endangered Taxa which is not critically endangered and it is facing a very high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
V	Vulnerable Taxa which is not critically endangered or endangered and is facing a high risk of extinction in the wild in the medium-term future, as determined in accordance with the prescribed criteria.
CD	Conservation Dependent Taxa which at a particular time if, at that time, the species is the focus of a specific conservation program, the cessation of which would result in the species becoming vulnerable, endangered or critically endangered within a period of 5 years.

Source: *Environment Protection and Biodiversity Conservation Act 1999*

A3: Definitions of Threatened Ecological Communities

Presumed Totally Destroyed (PD)

An ecological community will be listed as presumed totally destroyed if there are no recent records of the community being extant **and either** of the following applies (A or B);

- A) Records within the last 50 years have not been confirmed despite thorough searches or known or likely habitats **or**
- B) All occurrences recorded within the last 50 years have since been destroyed.

Critically Endangered (CR)

An ecological community will be listed as **Critically Endangered** when it has been adequately surveyed and is found to be facing an extremely high risk of total destruction in the immediate future. This will be determined on the basis of the best available information, by it meeting **any one or more** of the following criteria (A, B or C):

- A) The estimated geographic range, and/or total area occupied, and/or number of discrete occurrences since European settlement have been reduced by at least 90% and **either or both** of the following apply (i or ii)
 - i) geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is imminent (within approximately 5 years)
 - ii) modification throughout its range is continuing such that in the immediate future (within approximately 5 years) the community is unlikely to be capable of being substantially rehabilitated.
- B) Current distribution is limited, and **one or more** of the following apply (i, ii or iii):
 - i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the immediate future (within approximately 5 years)
 - ii) there are very few occurrences, each of which is small and/or isolated and extremely vulnerable to known threatening processes
 - iii) there may be many occurrences but total area is very small and each occurrence is small and/or isolated and extremely vulnerable to known threatening processes
- C) The ecological community exists only as highly modified occurrences which may be capable of being rehabilitated if such work begins in the immediate future (within approximately 5 years)

Endangered (EN)

An ecological community will be listed as **Endangered** when it has been adequately surveyed and is not Critically Endangered but is facing a very high risk of total destruction in the near future. This will be determined on the basis of the best available information, by it meeting **any one or more** of the following criteria (A, B or C):

- A) The estimated geographic range, and/or total area occupied, and/or number of discrete occurrences since European settlement have been reduced by at least 70% and **either or both** of the following apply (i or ii)
 - i) geographic range, and/or total area occupied and/or number of discrete occurrences are continuing to decline such that total destruction of the community is likely in the short term (within approximately 10 years)
 - ii) modification throughout its range is continuing such that in the short term future (within approximately 10 years) the community is unlikely to be capable of being substantially restored or rehabilitated.
- B) Current distribution is limited, and **one or more** of the following apply (i, ii or iii):
 - i) geographic range and/or number of discrete occurrences, and/or area occupied is highly restricted and the community is currently subject to known threatening processes which are likely to result in total destruction throughout its range in the short term future (within approximately 10 years)
 - ii) there are very few occurrences, each of which is small and/or isolated and extremely vulnerable to known threatening processes
 - iii) there may be many occurrences but total area is very small and each occurrence is small and/or isolated and extremely vulnerable to known threatening processes
- C) The ecological community exists only as highly modified occurrences which may be capable of being rehabilitated if such work begins in the short term future (within approximately 10 years).

Vulnerable (VU)

An ecological community will be listed as **Vulnerable** when it has been adequately surveyed and is not Critically Endangered or Endangered but is facing a high risk of total destruction in the medium to long term future. This will be determined on the basis of the best available information, by it meeting **any one or more** of the following criteria (A, B or C):

- A) The ecological community exists largely as modified occurrences which are likely to be capable of being substantially restored or rehabilitated.
- B) The ecological community can be modified or destroyed and would be vulnerable to threatening processes, is restricted in area and/or range and/or is only found at a few locations.
- C) The ecological community may still be widespread but is believed likely to move into a category of higher threat in the medium to long term future because of existing or impending threatening processes.

Source: Department of Environment and Conservation (2009). *Definitions, Categories and Criteria for Threatened and Priority Ecological Communities*. Department of Environment and Conservation, Perth, Western Australia. Online: www.naturebase.net/

A4: Definitions of Priority Ecological Communities

Possible threatened ecological communities that do not meet survey criteria or that are not adequately defined are added to the Priority Ecological Community Lists under Priorities 1, 2 and 3. These three categories are ranked in order of priority for survey and/or definition of the community, and evaluation of conservation status, so that consideration can be given to their declaration as threatened ecological communities. Ecological Communities that are adequately known, and are rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list, are placed in Priority 4. These ecological communities require regular monitoring. Conservation Dependent ecological communities are placed in Priority 5.

Priority One: Poorly known ecological communities Ecological communities with apparently few, small occurrences, all or most not actively managed for conservation (e.g. within agricultural or pastoral lands, urban areas, active mineral leases) and for which current threats exist. Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under immediate threat from known threatening processes across their range.

Priority Two: Poorly known ecological communities. Communities that are known from few small occurrences, all or most of which are actively managed for conservation (e.g. within national parks, conservation parks, nature reserves, State forest, unallocated Crown land, water reserves, etc.) and not under imminent threat of destruction or degradation.

Communities may be included if they are comparatively well known from one or more localities but do not meet adequacy of survey requirements, and/or are not well defined, and appear to be under threat from known threatening processes.

Priority Three: Poorly known ecological communities

- (i) Communities that are known from several to many occurrences, a significant number or area of which are not under threat of habitat destruction or degradation or;
- (ii) Communities known from a few widespread occurrences, which are either large or within significant remaining areas of habitat in which other occurrences may occur, much of it not under imminent threat, or;
- (iii) Communities made up of large, and/or widespread occurrences, that may or not be represented in the reserve system, but are under threat of modification across much of their range from processes such as grazing by domestic and/or feral stock, and inappropriate fire regimes.

Communities may be included if they are comparatively well known from several localities but do not meet adequacy of survey requirements and/or are not well defined, and known threatening processes exist that could affect them.

Priority Four: Ecological communities that are adequately known, rare but not threatened or meet criteria for Near Threatened, or that have been recently removed from the threatened list. These communities require regular monitoring.

- (a) Rare. Ecological communities known from few occurrences that are considered to have been adequately surveyed, or for which sufficient knowledge is available, and that are considered not currently threatened or in need of special protection, but could be if present circumstances change. These communities are usually represented on conservation lands.
- (b) Near Threatened. Ecological communities that are considered to have been adequately surveyed and that do not qualify for Conservation Dependent, but that are close to qualifying for Vulnerable.
- (c) Ecological communities that have been removed from the list of threatened communities during the past five years.

Priority Five: Conservation Dependent ecological communities. Ecological communities that are not threatened but are subject to a specific conservation program, the cessation of which would result in the community becoming threatened within five years.

Source: Department of Environment and Conservation (2009). *Definitions, Categories and Criteria for Threatened and Priority Ecological Communities*. Department of Environment and Conservation, Perth, Western Australia. Online: www.naturebase.net/

APPENDIX B

ENVIRONMENTAL WEEDS AND DECLARED PLANT CATEGORIES

APPENDIX B

ENVIRONMENTAL WEEDS AND DECLARED PLANT CATEGORIES

B1: Criteria used for Ranking Environmental Weeds

The Environmental Weed Strategy for Western Australia (CALM 1999) contains criteria for the assessment and ranking of weeds in terms of their environmental impact on biodiversity. These criteria are as follows:

- **Invasiveness** – ability to invade bushland in good to excellent condition or ability to invade waterways. (Score as yes or no).
- **Distribution** – wide current or potential distribution including consideration of known history of wide spread distribution elsewhere in the world. (Score as yes or no).
- **Environmental Impacts** – ability to change the structure, composition and function of ecosystems. In particular an ability to form a monoculture in a vegetation community. (Score as yes or no).

The rating of each weed is determined by the following scoring system:

- **High** - a weed species would have to score yes for all three criteria. Rating a weed species as high would indicate prioritising this weed for control and/or research i.e. prioritising funding to it.
- **Moderate** - a weed species would have to score yes for two of the above criteria. Rating a weed species as moderate would indicate that control or research effort should be directed to it if funds are available, however it should be monitored (possibly a reasonably high level of monitoring).
- **Mild** – a weed species scoring one of the criteria. A mild rating would indicate monitoring of the weed and control where appropriate.
- **Low** – a weed species would score none of the criteria. A low ranking would mean that this species would require a low level of monitoring.

Source: Department of Conservation and Land Management (1999). *Environmental Weed Strategy for Western Australia*. Department of Conservation and Land Management, Perth, Western Australia.

B2: Standard Meanings of Declared Plant Categories

P1

Prohibits movement.

The movement of plants or their seeds is prohibited within the State.

This prohibits the movement of contaminated machinery and produce including livestock and fodder.

P2

Aim is to eradicate infestation.

Treat all plants to destroy and prevent propagation each year until no plants remain. The infested area must be managed in such a way that prevents the spread of seed or plant parts on or in livestock, fodder, grain, vehicles and/or machinery.

P3

Aims to control infestation by reducing area and/or density of infestation.

The infested area must be managed in such a way that prevents the spread of seed or plant parts within and from the property on or in livestock, fodder, grain, vehicles and/or machinery.

Treat to destroy and prevent seed set all plants:

- * Within 50m inside of the boundaries of the infestation;
- * within 50m of roads and high water mark on waterways;
- * within 50m of sheds, stock yards and houses.

Treatment must be done prior to seed set each year.

Properties with less than 20ha of infestation must treat the entire infestation.

Additional areas may be ordered to be treated.

P4

Aims to prevent infestation spreading beyond existing boundaries of infestation

The infested area must be managed in such a way that prevents the spread of seed or plant parts within and from the property on or in livestock, fodder, grain, vehicles and/or machinery.

Treat to destroy and prevent seed set all plants:

- * within 50m inside of the boundaries of the infested property for one-leaf and 20m for two-leaf;
- * within 50m of roads and high water mark on waterways;
- * within 50m of sheds, stock yards and houses.

Treatment must be done prior to seed set each year. Properties with less than 20ha of infestation must treat the entire infestation.

Additional areas may be ordered to be treated.

Special considerations.

In the case of P4 infestations where they continue across property boundaries there is no requirement to treat the relevant part of the property boundaries as long as the boundaries of the infestation as a whole are treated. There must be agreement between neighbours in relation to the treatment of these areas.

P5

Aims to control infestations on public lands.

Source: Department of Agriculture and Food (2008). *List of Declared Plants*. Department of Agriculture and Food, Western Australia. Online: <http://www.agric.wa.gov.au/>.

APPENDIX C

DEC DATABASE SEARCH RESULTS

APPENDIX C

DEPARTMENT OF ENVIRONMENT AND CONSERVATION DATABASE
SEARCH RESULTS

C1: Declared Rare and Priority Flora

FAMILY	TAXA	Conservation Status Code	
		STATE	FEDERAL
ORCHIDACEAE	<i>Caladenia huegelii</i>	R	ENDANGERED
ORCHIDACEAE	<i>Drakaea elastica</i>	R	ENDANGERED
MYRTACEAE	<i>Verticordia plumosa</i> var. <i>ananeotes</i>	R	ENDANGERED
ORCHIDACEAE	<i>Diuris micrantha</i>	R	VULNERABLE
MYRTACEAE	<i>Eremaea asterocarpa</i> subsp. <i>brachyclada</i>	P1	Not Listed
MIMOSACEAE	<i>Acacia benthamii</i>	P2	Not Listed
PAPILIONACEAE	<i>Aotus cordifolia</i>	P3	Not Listed
EUPHORBIACEAE	<i>Beyeria cinerea</i> subsp. <i>cinerea</i>	P3	Not Listed
CYPERACEAE	<i>Cyathochaeta teretifolia</i>	P3	Not Listed
PAPILIONACEAE	<i>Dillwynia dillwynioides</i>	P3	Not Listed
GOODENIACEAE	<i>Goodenia filiformis</i>	P3	Not Listed
PAPILIONACEAE	<i>Jacksonia gracillima</i>	P3	Not Listed
CYPERACEAE	<i>Schoenus capillifolius</i>	P3	Not Listed
STYLIDIACEAE	<i>Stylidium longitubum</i>	P3	Not Listed
APONOGETONACEAE	<i>Aponogeton hexatpalus</i>	P4	Not Listed
SAPINDACEAE	<i>Dodonaea hackettiana</i>	P4	Not Listed
BRASSICACEAE	<i>Lepidium puberulum</i>	P4	Not Listed
PAPILIONACEAE	<i>Jacksonia sericea</i>	P4	Not Listed
STYLIDIACEAE	<i>Stylidium ireneae</i>	P4	Not Listed

C2: Threatened and Priority Ecological Communities

- SCP19a** Sedgelands in Holocene dune swales of the southern Swan Coastal Plain (listed as Critically Endangered under the federal legislation)
- SCP19b** Woodlands over sedgelands in Holocene dune swales of the southern Swan (listed as Critically Endangered under the federal legislation)
- Richmond-microbial** Thrombolite like microbialite community of coastal freshwater lakes (listed as Critically Endangered under the federal legislation)
- SCP30a** *Callitris preissii* (or *Melaleuca lanceolata*) forests and woodlands (listed as Vulnerable under the federal legislation)

APPENDIX D

FLORA SPECIES LIST

APPENDIX D

MATRIX OF SPECIES FOUND AT EACH SITE

Taxa	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13	Q14	Q15	Q16	OPP COLLS	Additional species recorded by Bennett Consulting Pty Ltd (2005)
<i>Acacia ? cyclops</i>		1%																
<i>Acacia cochlearis</i>		5%																
<i>Acacia cyclops</i>									+		2%	1.5%		2%				
<i>Acacia lasiocarpa</i> var. <i>lasiocarpa</i>																		+
<i>Acacia pulchella</i>								+										
<i>Acacia rostellifera</i>	2%					nc		90%		5%			75%			90%		
<i>Acacia saligna</i>							2%									1.5%		
<i>Acanthocarpus preissii</i>	nc		1%	3%		10%	10%	3%			3%			1%		5%		
<i>Agonis flexuosa</i>									75%		40%							
<i>Allocasuarina lehmanniana</i>																		+
<i>Alyxia buxifolia</i>			10%						15%	20%		1%		1.5%				
* <i>Lysimachia arvensis</i>				+				+			1%							
* <i>Arctotheca calendula</i>		+																
* <i>Asparagus asparagoides</i>								nc										
* <i>Asphodelus fistulosus</i>																		+
<i>Atriplex isatidea</i>																		+
* <i>Avena barbata</i>	+	10%		+			1%			8%	2%		2%					
* <i>Bromus diandrus</i>	1%	1%	+		nc		+	+						5%	1.5%			
* <i>Cakile maritima</i>										1%								
<i>Callitris preissii</i>		2%									25%							
<i>Calothamnus quadrifidus</i>								nc										
* <i>Carpobrotus edulis</i>			nc															
<i>Cassytha racemosa</i>						nc							6%	3%				
<i>Clematis linearifolia</i>	+		1%								+							
<i>Conostylis candicans</i>							+	nc								+		
* <i>Conyza parva</i>																		+
* <i>Crassula glomerata</i>						+		+						1%	1.5%			
* <i>Cuscuta epithymum</i>			+															
* <i>Cynodon dactylon</i>													1.5%					
* <i>Cyperus congestus</i>																		+
<i>Desmocladius flexuosus</i>								2%										
<i>Dianella revoluta</i>											+							
<i>Diplolaena dampieri</i>																	+	
* <i>Ehrharta calycina</i>	+								2%				5%					
* <i>Ehrharta longiflora</i>	+																	
* <i>Eragrostis curvula</i>																3%		
<i>Eremophila glabra</i> subsp. <i>albicans</i>	nc	nc														1.5%		
* <i>Erodium</i> sp.																		+
<i>Eucalyptus gomphocephala</i>										7%	60%							
* <i>Eucalyptus utilis</i>																		+
* <i>Euphorbia paralias</i>					1%													
* <i>Euphorbia peplus</i>											6%							
* <i>Euphorbia terracina</i>	nc	2%	+	+		+	2%	1%		2%	1%		1%	1%		4%		
<i>Exocarpos sparteus</i>						3%												
<i>Ficinia nodosa</i>					+													
* <i>Ficus carica</i>																		+
* <i>Foeniculum vulgare</i>								nc										
<i>Frankenia pauciflora</i>			nc		1%													
* <i>Fumaria capreolata</i>	60%							nc		8%								
* <i>Gazania linearis</i>																		+

APPENDIX D

MATRIX OF SPECIES FOUND AT EACH SITE

Taxa	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13	Q14	Q15	Q16	OPP COLLS	Additional species recorded by Bennett Consulting Pty Ltd (2005)
<i>*Geranium molle</i>											+							
<i>Hardenbergia comptoniana</i>	15%			+		+					nc							
<i>Hemiandra pungens</i>																		+
<i>Hibbertia cuneiformis</i>																		0
<i>*Hyparrhenia hirta</i>							60%											
<i>Jacksonia furcellata</i>							nc											
<i>*Lagurus ovatus</i>		+		nc			1%	2%			1%			3%		5%		
<i>Lepidosperma gladiatum</i>	35%	nc	15%	50%		3%			50%	15%								
<i>Lepidosperma</i> sp. Coastal Dune (R. J. Cranfield 9963)								nc										
<i>*Leptospermum laevigatum</i>																		+
<i>Leucopogon parviflorus</i>											+					+		
<i>*Lolium rigidum</i>	10%	2%	+	1%	nc	+	1%	1%	1%	10%	4%		5%	1%	3%	3%		
<i>Lomandra maritima</i>								+								1%		
<i>*Malva dendromorpha</i>																		+
<i>*Melaleuca diosmifolia</i>																		+
<i>Melaleuca huegelii</i> subsp. <i>huegelii</i>		2%																
<i>Melaleuca lanceolata</i>		16%										90%						
<i>*Melaleuca nesophila</i>																		+
<i>*Melilotus indicus</i>			nc	10%	+													
<i>Myporum insulare</i>																		+
<i>*Olea europaea</i>											1%							
<i>Olearia axillaris</i>		2%	nc		+	nc			nc						15%			
<i>Ozothamnus cordatus</i>						4%												
<i>*Pelargonium capitatum</i>			2%	1%			1%	1%		2%	2%			6%	5%	2%		
<i>*Pennisetum clandestinum</i>									2%									
<i>Phyllanthus calycinus</i>								10%								1%		
<i>Pittosporum ligustrifolium</i>				nc														
<i>*Plantago lanceolata</i>							nc											
POACEAE sp.				3%		1%		1%										
<i>*Raphanus raphanistrum</i>																		+
<i>Rhagodia baccata</i>	2%	2%	5%	5%		+		nc	3%		2%	7%	8%	1.5%				
<i>*Rhamnus alaternus</i>											4%							
<i>*Romulea rosea</i>											nc							
<i>Salsola tragus</i>																		
<i>Scaevola crassifolia</i>		2%	nc	1%	+	20%		2%							1.5%			
<i>*Schinus terebinthifolius</i>							2%				3%							
<i>Senecio pinnatifolius</i>	+	3%	1%	+		1%		+										
<i>*Sonchus oleraceus</i>	nc		+						+				+	+				
<i>Spinifex hirsutus</i>																1%		
<i>Spinifex longifolius</i>						nc				2%				15%	10%			
<i>Spyridium globulosum</i>	nc	3%	10%					+			60%							
<i>*Symphyotrichum subulatum</i>																		+
<i>Templetonia retusa</i>		1%																
<i>*Tetragonia decumbens</i>			10%			5%			3%	2%		1%		+	+			
<i>Threlkeldia diffusa</i>		+	1%															
<i>*Trachyandra divaricata</i>			30%	2%	nc	1%	+			+			1.5%	6%	20%			
<i>*Urospermum picroides</i>	10%																	
<i>*Ursinia anthemoides</i>																		+
<i>*Westringia fruticosa</i>																		+

APPENDIX E

FLORA SURVEY FIELD DATASHEETS AND SITE PHOTOGRAPHS

APPENDIX E

FLORA SURVEY FIELD DATA SHEETS AND PHOTOGRAPHS

Site Q01

Described by NP

Location Point Peron.

MGA Zone 50 376614 mE 6428940 mN

Habitat Sloping fore dune on dark grey to black sand

Soil Grey to black sand.

Rock Type N/A.

Vegetation SCP29b

Veg Condition Very Good.

Fire Age Old.

Notes Aspect: South-East.
Bare ground: 0%.
Litter cover: 0% Logs 0% Twigs 0% Lvs.
Disturbance type: Weeds.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia rostellifera</i>	2%	1-2m	Q1-07
<i>Acanthocarpus preissii</i>		0.25m	Associated.
* <i>Avena barbata</i>	+	1.5m	Q1-11
* <i>Bromus diandrus</i>	1%	0.5m	Q1-03
<i>Clematis linearifolia</i>	+	C	Q1-08
* <i>Ehrharta calycina</i>	+	0.5m	Q1-04
* <i>Ehrharta longiflora</i>	+	0.3m	Q1-09
<i>Eremophila glabra</i> subsp. <i>albicans</i>		0.4m	Q1-13 Associated.
* <i>Euphorbia terracina</i>		0.3m	Associated.
* <i>Fumaria capreolata</i>	60%	1m	
<i>Hardenbergia comptoniana</i>	15%	C	Q1-02
<i>Lepidosperma gladiatum</i>	35%	1.2m	Q1-01
* <i>Lolium rigidum</i>	10%	0.4m	Q4-02
<i>Rhagodia baccata</i>	2%	0.6m	Q1-05
<i>Senecio pinnatifolius</i>	+	0.25m	Q1-10
* <i>Sonchus oleraceus</i>		0.3m	Q1-14 Associated.
<i>Spyridium globulosum</i>		2.1m	Associated.

Site Q02

Described by NP

Location Point Peron.
MGA Zone 50 376415 mE 6428720 mN
Habitat Shrubland.
Soil Dark yellow sand.
Rock Type Small amounts of limestone.
Vegetation SCP30a
Veg Condition Good to Very Good.
Fire Age Old.
Notes Aspect: North-West.
 Bare ground: 10%.
 Litter cover: 5% Logs 1% Twigs +% Lvs.
 Disturbance type: Weeds; surrounding
 paths/roads/carpark/lookout.
 Notes: Recent rehabilitation on bare areas.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia ? cyclops</i>	1%	0.9m	Q2-05
<i>Acacia cochlearis</i>	5%	1m	Q2-04
* <i>Arctotheca calendula</i>	+	0.15m	
* <i>Avena barbata</i>	10%	0.6m	Q2-03
* <i>Bromus diandrus</i>	1%	0.2m	Q1-03
<i>Callitris preissii</i>	2%	1.5m	Q2-07
<i>Eremophila glabra</i> subsp. <i>albicans</i>		1m	Q1-13 Associated.
* <i>Euphorbia terracina</i>	2%	0.2m	
* <i>Lagurus ovatus</i>	+	0.4m	Q2-11
<i>Lepidosperma gladiatum</i>		1m	Q1-01 Associated.
* <i>Lolium rigidum</i>	2%	0.2m	Q4-02
<i>Melaleuca huegelii</i> subsp. <i>huegelii</i>	2%	1.2m	Q2-08
<i>Melaleuca lanceolata</i>	16%	1.2m	Q2-06
<i>Olearia axillaris</i>	2%	1.2m	
<i>Rhagodia baccata</i>	2%	0.7m	Q1-05
<i>Scaevola crassifolia</i>	2%	0.35m	Q2-12
<i>Senecio pinnatifolius</i>	3%	0.3m	Q1-10
<i>Spyridium globulosum</i>	3%	0.8m	
<i>Templetonia retusa</i>	1%	1.5m	Q2-13
<i>Threlkeldia diffusa</i>	+	0.1m	Q2-02
* <i>Urospermum picroides</i>	10%	0.15m	Q2-01

Site Q03

Described by NP

Location Point Peron.
MGA Zone 50 376277 mE 6428772 mN
Habitat Dune slope to limestone knoll.
Soil White/Yellow sand.
Rock Type Limestone.
Vegetation SCP16 and SCP29a
Veg Condition Good.
Fire Age Old.
Notes Aspect: West.
 Bare ground: 50%.
 Litter cover: 0% Logs; 0% Twigs; 1% Lvs.
 Disturbance type: Weeds, walk trails, rubbish.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acanthocarpus preissii</i>	1%	0.2m	
<i>Alyxia buxifolia</i>	10%	1m	Q3-01
* <i>Bromus diandrus</i>	+	0.2m	Q1-03
* <i>Carpobrotus edulis</i>		0.1m	Associated.
<i>Clematis linearifolia</i>	1%	C	Q1-08
* <i>Cuscuta epithymum</i>	+	<0.1m	Q3-06
* <i>Euphorbia terracina</i>	+	0.1m	
<i>Frankenia pauciflora</i>		0.3m	Q3-08 Associated.
<i>Lepidosperma gladiatum</i>	15%	1m	Q1-01
* <i>Lolium rigidum</i>	+	0.25m	Q4-02
* <i>Melilotus indicus</i>		0.2m	Q3-07 Associated.
<i>Olearia axillaris</i>		0.4m	Associated.
* <i>Pelargonium capitatum</i>	2%	0.3m	Q3-05
<i>Rhagodia baccata</i>	5%	0.9m	Q3-02
<i>Scaevola crassifolia</i>		1m	Q2-12 Associated.
<i>Senecio pinnatifolius</i>	1%	0.3m	Q1-10
* <i>Sonchus oleraceus</i>	+	0.1m	Q3-04
<i>Spyridium globulosum</i>	10%	0.9m	
<i>Tetragonia decumbens</i>	10%	0.4m	Q3-03
<i>Threlkeldia diffusa</i>	1%	0.1m	Q2-02
* <i>Trachyandra divaricata</i>	30%	0.3m	

Site Q04

Described by NP

Location Point Peron.
MGA Zone 50 376273 mE 6429363 mN
Habitat Sloping fore dune
Soil Grey Sand.
Rock Type N/A.
Vegetation S14
Veg Condition Very Good.
Fire Age Old.
Notes Aspect: South.
 Bare ground: 0%.
 Litter cover: 0% Logs; 0% Twigs; 0% Lvs.
 Disturbance type: Weeds.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acanthocarpus preissii</i>	3%	0.3m	
* <i>Lysimachia arvensis</i>	+	<0.1m	
* <i>Avena barbata</i>	+	0.5m	Q4-04
* <i>Euphorbia terracina</i>	+	0.4m	
<i>Hardenbergia comptoniana</i>	+	C	Q4-05
* <i>Lagurus ovatus</i>		0.2m	Q2-11
<i>Lepidosperma gladiatum</i>	50%	0.9m	Q1-01
* <i>Lolium rigidum</i>	1%	0.3m	Q4-02
* <i>Melilotus indicus</i>	10%	0.2m	Q3-07
* <i>Pelargonium capitatum</i>	1%	0.2m	Q3-05
<i>Pittosporum ligustrifolium</i>		1m	Q4-06
POACEAE sp.	3%	1m	Q4-01
<i>Rhagodia baccata</i>	5%	0.3m	Q4-03
<i>Scaevola crassifolia</i>	1%	0.9m	Q2-12
<i>Senecio pinnatifolius</i>	+	0.3m	Q1-10
* <i>Trachyandra divaricata</i>	2%	0.4m	

Site Q05**Described by NP**

Location Point Peron.
MGA Zone 50 376271 mE 6429416 mN
Habitat Ironstone outcrop of cliff/knoll.
Soil Red/Yellow sand.
Rock Type Limestone.
Vegetation SCP16 and SCP29a
Veg Condition Completely Degraded.
Fire Age Old.
Notes Aspect: North.
 Bare ground: 80%.
 Litter cover: 0% Logs; 0% Twigs; 0% Lvs.

**SPECIES LIST:**

Name	Cover	Height	Specimen	Notes
* <i>Bromus diandrus</i>		0.2m	Q1-03	Associated.
* <i>Euphorbia paralias</i>	1%	0.2m	Q5-01	
<i>Ficinia nodosa</i>	+	0.9m	Q5-02	
<i>Frankenia pauciflora</i>	1%	0.2m	Q3-08	
* <i>Lolium rigidum</i>		0.3m	Q4-02	Associated.
* <i>Melilotus indicus</i>	+	0.1m	Q3-07	
<i>Olearia axillaris</i>	+	0.2m		
<i>Scaevola crassifolia</i>	+	<0.1m	Q2-12	
* <i>Trachyandra divaricata</i>		0.1m		Associated.

Site Q06

Described by NP

Location Point Peron.
MGA Zone 50 376341 mE 6428938 mN
Habitat Fore dune
Soil Yellow/Brown sand.
Rock Type N/A.
Vegetation S13
Veg Condition Very Good to Excellent.
Fire Age Old.
Notes Aspect: West.
 Bare ground: 1%.
 Litter cover: 0% Logs; +% Twigs; + %Lvs.
 Disturbance type: Weeds.



SPECIES LIST:

Name	Cover	Height	Specimen	Notes
<i>Acacia rostellifera</i>		0.3m	Q6-06	Associated.
<i>Acanthocarpus preissii</i>	10%	0.3m		
<i>Cassyltha racemosa</i>		C	Q6-05	Associated.
* <i>Crassula glomerata</i>	+	<0.1m	Q6-02	
* <i>Euphorbia terracina</i>	+	0.2m		
<i>Exocarpos sparteus</i>	3%	2m	Q6-03	
<i>Hardenbergia comptoniana</i>	+	C	Q1-02	
<i>Lepidosperma gladiatum</i>	3%	0.9m	Q1-01	
* <i>Lolium rigidum</i>	+	0.25m	Q4-02	
<i>Olearia axillaris</i>		1.5m		Associated.
<i>Ozothamnus cordatus</i>	4%	1m	Q6-01	
POACEAE sp.	1%	0.5m	Q4-01	
<i>Rhagodia baccata</i>	+	0.2m	Q1-05	
<i>Scaevola crassifolia</i>	20%	0.5m	Q2-12	
<i>Senecio pinnatifolius</i>	1%	0.3m	Q1-10	
<i>Spinifex longifolius</i>		0.45m	Q6-04	Associated.
* <i>Tetragonia decumbens</i>	5%	0.2m	Q3-03	
* <i>Trachyandra divaricata</i>	1%	0.3m		

Site Q07

Described by NP

Location Point Peron
MGA Zone 50 378489 mE 6428101 mN
Habitat Weed dominated shrubland.
Soil Yellow to Grey sand.
Rock Type N/A.
Vegetation S15
Veg Condition Completely Degraded.
Fire Age Old.
Notes Aspect: N/A.
 Bare ground: 10%.
 Litter cover: 0% Logs; 0% Twigs; 0% Lvs.
 Disturbance type: Weeds, rubbish.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia saligna</i>	2%	1.1m	Q7-02
<i>Acanthocarpus preissii</i>	10%	0.4m	
* <i>Avena barbata</i>	1%	0.7m	Q1-11
* <i>Bromus diandrus</i>	+	0.25m	Q1-03
<i>Conostylis candicans</i>	+	0.3m	Q7-04
* <i>Euphorbia terracina</i>	2%	0.4m	
* <i>Foeniculum vulgare</i>		2m	Associated.
* <i>Fumaria capreolata</i>		0.3m	Associated.
* <i>Hyparrhenia hirta</i>	60%	1.5m	Q7-01
<i>Jacksonia furcellata</i>		2.2m	Associated.
* <i>Lagurus ovatus</i>	1%	0.3m	Q2-11
* <i>Lolium rigidum</i>	1%	0.3m	Q4-02
* <i>Pelargonium capitatum</i>	1%	0.3m	Q3-05
* <i>Plantago lanceolata</i>		0.5m	Q7-03
* <i>Schinus terebinthifolius</i>	2%	2m	Associated.
* <i>Trachyandra divaricata</i>	+	0.3m	

Site Q08

Described by NP

Location Point Peron.
MGA Zone 50 377879 mE 6428061 mN
Habitat Shrubland dominated by *Acacia rostellifera*.
Soil Yellow to light brown/red sand.
Rock Type N/A.
Vegetation SCP29b
Veg Condition Excellent.
Fire Age Old.
Notes Aspect: North-West.
 Bare ground: 1%.
 Litter cover: 3% Logs; 1% Twigs; 1% Lvs.
 Disturbance type: Weeds, rabbits.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia pulchella</i>	+	0.2m	
<i>Acacia rostellifera</i>	90%	2m	Q8-01
<i>Acanthocarpus preissii</i>	3%	0.3m	
* <i>Lysimachia arvensis</i>	+	0.1m	
* <i>Asparagus asparagoides</i>		0.1m	Associated.
* <i>Bromus diandrus</i>	+	0.2m	Q1-03
<i>Calothamnus quadrifidus</i>		1.2m	Q8-06 Associated.
<i>Conostylis candicans</i>		0.2m	Q8-07 Associated.
* <i>Crassula glomerata</i>	+	<0.1m	Q6-02
<i>Desmocladius flexuosus</i>	2%	0.15m	Q8-09
* <i>Euphorbia terracina</i>	1%	0.15m	
* <i>Lagurus ovatus</i>	2%	0.2m	Q8-02
<i>Lepidosperma</i> sp. Coastal Dune (R. J. Cranfield 9963)		0.15m	Q8.11 Associated
* <i>Lolium rigidum</i>		1%	0.4m Q4-02
<i>Lomandra maritima</i>	+	0.25m	Q8-08
* <i>Pelargonium capitatum</i>	1%	0.3m	Q3-05
<i>Phyllanthus calycinus</i>	10%	0.3m	
POACEAE sp.		1m	Q4-01 Associated.
POACEAE sp.	1%	0.4m	Q8-03
<i>Rhagodia baccata</i>		1m	Q8-05 Associated.
<i>Scaevola crassifolia</i>	2%	0.2m	Q2-12
<i>Senecio pinnatifolius</i>	+	0.3m	Q8-10
<i>Spyridium globulosum</i>	+	0.2m	

Site Q09

Described by NP

Location Point Peron.
MGA Zone 50 376977 mE 6428639 mN
Habitat Woodland over shrubs and sedges.
Soil Grey sand.
Rock Type N/A.
Vegetation SCP30b
Veg Condition Good.
Fire Age Very Old.
Notes Aspect: N/A.
 Bare ground: 0%.
 Litter cover: 0% Logs, 2% Twigs; 3% Lvs.
 Disturbance type: Weeds.
 Notes: Road within 5m of community.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia cyclops</i>	+	0.45m	NW4
<i>Agonis flexuosa</i>	75%	9m	
<i>Alyxia buxifolia</i>	15%	1.1m	NW1
* <i>Ehrharta calycina</i>	2%	0.6m	
<i>Lepidosperma gladiatum</i>	50%	1.2m	
* <i>Lolium rigidum</i>	1%	0.45m	NW3
<i>Olearia axillaris</i>			Opportunistic.
* <i>Pennisetum clandestinum</i>	2%	0.45m	
<i>Rhagodia baccata</i>	3%	1m	NW5
* <i>Sonchus oleraceus</i>	+	0.3m	
* <i>Tetragonia decumbens</i>	3%	1m	NW2

Site Q10

Described by NW

Location Point Peron.
MGA Zone 50 376963 mE 6428593 mN
Habitat Dunes.
Soil Sand (dune).
Rock Type N/A.
Vegetation SCP30b
Veg Condition Good.
Fire Age Old
Notes Aspect: East.
 Bare ground: 2%.
 Litter cover: 1% Logs 1% Twigs 2% Lvs.
 Disturbance type: Weeds.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia rostellifera</i>	5%	1.2m	NW6
<i>Alyxia buxifolia</i>	20%	1.3m	
* <i>Avena barbata</i>	8%	0.6m	
* <i>Cakile maritima</i>	1%	0.35m	NW7
<i>Eucalyptus gomphocephala</i>	7%	11m	
* <i>Euphorbia terracina</i>	2%	0.35m	
* <i>Fumaria capreolata</i>	8%	0.4m	
<i>Lepidosperma gladiatum</i>	15%	1.2m	
* <i>Lolium rigidum</i>	10%	0.5m	NW3
* <i>Pelargonium capitatum</i>	2%	0.45m	
<i>Spinifex longifolius</i>	2%	0.85m	NW8
* <i>Tetragonia decumbens</i>	2%	0.35m	NW2
* <i>Trachyandra divaricata</i>	+	0.4m	

Site Q11

Described by NW

Location Point Peron.
MGA Zone 50 378408 mE 427831 mN
Habitat Woodland.
Soil Grey sand.
Rock Type N/A.
Vegetation SCP30a
Veg Condition Good.
Fire Age Very Old.
Notes Aspect: N/A.
 Bare ground: 0%.
 Litter cover: 0% logs; 6% Twigs; 20% Lvs.
 Disturbance type: Weeds, areas of clearing nearby, tracks.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia cyclops</i>	2%	120m	
<i>Acanthocarpus preissii</i>	3%	0.35m	
<i>Agonis flexuosa</i>	40%	8m	
* <i>Lysimachia arvensis</i>	1%	0.05m	
* <i>Avena barbata</i>	2%	0.4m	
<i>Callitris preissii</i>	25%	2.3m	
<i>Clematis linearifolia</i>	+	1.6m	
<i>Dianella revoluta</i>	+	0.45m	
<i>Eucalyptus gomphocephala</i>	60%	16m	
* <i>Euphorbia peplus</i>	6%	0.2m	
* <i>Euphorbia terracina</i>	1%	0.4m	
* <i>Geranium molle</i>	+	0.1m	
<i>Hardenbergia comptoniana</i>			
* <i>Lagurus ovatus</i>	1%	0.25m	
<i>Leucopogon parviflorus</i>	+	C	
* <i>Lolium rigidum</i>	4%	0.5m	NW3
* <i>Olea europaea</i>	1%	0.1m	NW10
* <i>Pelargonium capitatum</i>	2%	1m	
<i>Rhagodia baccata</i>	2%	0.1m	
* <i>Rhamnus alaternus</i>	4%	2.2m	NW11
* <i>Romulea rosea</i>		1.8m	
* <i>Schinus terebinthifolius</i>	3%	1.5m	
<i>Spyridium globulosum</i>	60%	1.7m	NW9

Site Q12

Described by NW

Location Point Peron.
MGA Zone 50 376890 mE 6428951 mN
Habitat Thick tall shrubland with no understorey.
Soil Sand - coastal.
Rock Type N/A.
Vegetation SCP30a
Veg Condition Good to Degraded.
Fire Age Very Old.
Notes Aspect: N/A.
 Bare ground: 4%.
 Litter cover: 2% Logs; 5% Twigs; 20% Lvs.
 Notes: On Western edge of school oval and
 very thin strip <10m; all the *Callitis preissii* are dead.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia cyclops</i>	1.5%	1m	
<i>Alyxia buxifolia</i>	1%	1.1m	
<i>Melaleuca lanceolata</i>	90%	9m	
<i>Rhagodia baccata</i>	7%	1.2m	
* <i>Tetragonia decumbens</i>	1%	0.4m	

Site Q13

Described by NW

Location Point Peron.
MGA Zone 50 376982 mE 6428910 mN
Habitat Tall Shrubland.
Soil Sand - coastal dunes.
Rock Type N/A.
Vegetation SCP29a
Veg Condition Very Good to Good.
Fire Age Very Old.
Notes Aspect: N/A.
 Bare ground: 3%.
 Litter cover: 0% Logs; 4% Twigs; 25% Lvs.
 Disturbance type: Weeds.
 Notes: Acacia form a dense canopy.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia rostellifera</i>	75%	3m	
* <i>Avena barbata</i>	2%	0.45m	
<i>Cassutha racemosa</i>	6%	C	
* <i>Cynodon dactylon</i>	1.5%	0.15m	
* <i>Ehrharta calycina</i>	5%	0.45m	
* <i>Euphorbia terracina</i>	1%	0.4m	
* <i>Lolium rigidum</i>	5%	0.35m	
<i>Rhagodia baccata</i>	8%	1.1m	
* <i>Sonchus oleraceus</i>	+	0.25m	
* <i>Trachyandra divaricata</i>	1.5%	0.35m	

Site Q14

Described by NW

Location Point Peron
MGA Zone 50 376969 mE 6428955 mN
Habitat Coastal Dunes.
Soil Dunal sand.
Rock Type N/A.
Vegetation S14
Veg Condition Good to Degraded.
Fire Age Very Old.
Notes Aspect: N/A.
 Bare ground: 20%.
 Litter cover: 0% Logs; 2% Twigs; 1% Lvs.
 Disturbance type: Weeds, wind erosion.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia cyclops</i>	2%	1.3m	
<i>Acanthocarpus preissii</i>	1%	0.15m	
<i>Alyxia buxifolia</i>	1.5%	1m	NW1
* <i>Bromus diandrus</i>	5%	0.25m	NW13
<i>Cassytha racemosa</i>	3%	C	
* <i>Crassula glomerata</i>	1%	0.03m	NW12
* <i>Euphorbia terracina</i>	1%	0.35m	
* <i>Lagurus ovatus</i>	3%	0.3m	
* <i>Lolium rigidum</i>	1%	0.3m	
* <i>Pelargonium capitatum</i>	6%	0.5m	
<i>Rhagodia baccata</i>	1.5%	0.45m	
* <i>Sonchus oleraceus</i>	+	0.4m	
<i>Spinifex longifolius</i>	15%	0.5m	
* <i>Tetragonia decumbens</i>	+	0.15m	
* <i>Trachyandra divaricata</i>	6%	0.4m	

Site Q15

Described by NW

Location Point Peron.
MGA Zone 50 377207 mE 6428253 mN
Habitat
Soil Dune sand.
Rock Type N/A.
Vegetation S13
Veg Condition Very Good to Good.
Fire Age Very Old.
Notes Aspect: N/A - top of dune system.
 Bare ground: 25%.
 Litter cover: 0% Logs; 2% Twigs; 1% Lvs.
 Disturbance type: Weeds, erosion.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
* <i>Bromus diandrus</i>	1.5%	0.25m	NW13
* <i>Crassula glomerata</i>	1.5%	0.03m	NW12
* <i>Lolium rigidum</i>	3%	0.35m	
<i>Olearia axillaris</i>	15%	1.2m	
* <i>Pelargonium capitatum</i>	5%	0.45m	
<i>Scaevola crassifolia</i>	1.5%	0.45m	
<i>Spinifex hirsutus</i>	1%	0.7m	
<i>Spinifex longifolius</i>	10%	1m	
* <i>Tetragonia decumbens</i>	+	0.15m	
* <i>Trachyandra divaricata</i>	20%	0.35m	

Site Q16

Described by NW

Location Point Peron
MGA Zone 50 378172 mE 6427478 mN
Habitat Shrubland.
Soil Yellow/Brown sand.
Rock Type N/A.
Vegetation SCP29a
Veg Condition Good to Degraded.
Fire Age Moderate.
Notes Aspect: South-East.
 Bare ground: 3%.
 Litter cover: 0% Logs; 2% Twigs; 2% Lvs.
 Disturbance type: Weeds.



SPECIES LIST:

Name	Cover	Height	Specimen Notes
<i>Acacia rostellifera</i>	90%	3m	
<i>Acacia saligna</i>	1.5%	2.5m	
<i>Acanthocarpus preissii</i>	5%	0.4m	
<i>Conostylis candicans</i>	+	0.35m	
* <i>Eragrostis curvula</i>	3%	1.1m	
<i>Eremophila glabra</i> subsp. <i>albicans</i>	1.5%	0.65m	NW14
* <i>Euphorbia terracina</i>	4%	0.45m	
* <i>Lagurus ovatus</i>	5%	0.35m	
<i>Leucopogon parviflorus</i>	+	0.4m	
* <i>Lolium rigidum</i>	3%	0.3m	
<i>Lomandra maritima</i>	1%	0.35m	
* <i>Pelargonium capitatum</i>	2%	0.45m	
<i>Phyllanthus calycinus</i>	1%	0.4m	

APPENDIX F

BUSH FOREVER CONDITION SCALE

APPENDIX F

BUSH FOREVER CONDITION SCALE

Condition Scale Code	Condition Scale
P	Pristine (1) Pristine or nearly so, no obvious signs of disturbance
E	Excellent (2) Vegetation structure intact, disturbance affecting individual species and weeds are non-aggressive species.
VG	Very Good (3) Vegetation structure altered, obvious signs of disturbance. For example, disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.
G	Good (4) Vegetation structure significantly altered by very obvious signs of multiple disturbance. Retains basic vegetation structure or ability to regenerate it. For example, disturbance to vegetation structure caused by very frequent fires, the presence of some very aggressive weeds at high density, partial clearing, dieback and grazing.
D	Degraded (5) Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. For example, disturbance to vegetation structure caused by very frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.
CD	Completely Degraded (6) The structure of the vegetation is no longer intact and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees or shrubs.

Source: Government of Western Australia (2000). *Bush Forever Volume 2: Directory of Bush Forever Sites*. Department of Environmental Protection, Perth, Western Australia.

APPENDIX G

FLORISTICS DENDOGRAM

APPENDIX G FLORISTICS DENDROGRAM

